

THERMODYNAMICS OF MIXTURES CONTAINING AROMATIC ALCOHOLS. 1. LIQUID-LIQUID EQUILIBRIA FOR PHENYL METHANOL + n-ALKANE SYSTEMS

Journal:	<i>Journal of Chemical & Engineering Data</i>
Manuscript ID:	je-2012-00146k.R2
Manuscript Type:	Article
Date Submitted by the Author:	14-Mar-2012
Complete List of Authors:	Gonzalez, Juan; University of Valladolid, Física Aplicada Alonso-Tristán, Cristina; UNIVERSITY OF BURGOS, INGENIERÍA ELECTROMECÁNICA García de la Fuente, Isaias; UNiveridad de Valladolid, Física Aplicada Cobos, Jose; Universidad de Valladolid, Dpto Termodinamica y Fisica Aplicada

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45

**THERMODYNAMICS OF MIXTURES CONTAINING AROMATIC
ALCOHOLS. 1. LIQUID-LIQUID EQUILIBRIA FOR
(PHENYLMETHANOL + ALKANE) SYSTEMS**

JUAN ANTONIO GONZÁLEZ^(1,*), CRISTINA ALONSO-TRISTÁN⁽²⁾, ISAÍAS
GARCÍA DE LA FUENTE⁽¹⁾ AND JOSÉ CARLOS COBOS⁽¹⁾

⁽¹⁾ **G.E.T.E.F., Grupo Especializado en Termodinámica de Equilibrio entre Fases,**
Departamento de Física Aplicada, Facultad de Ciencias, Universidad de Valladolid, E-47071,
Valladolid (Spain).

⁽²⁾ Dpto. Ingeniería Electromecánica. Escuela Politécnica Superior. Avda. Cantabria s/n. 09006
Burgos, (Spain)

46 *corresponding author

47 E-mail: jagl@termo.uva.es
48
49
50
51
52
53
54
55
56
57
58
59
60

ABSTRACT

The liquid-liquid equilibrium (LLE) curves for (phenylmethanol + $\text{CH}_3(\text{CH}_2)_n\text{CH}_3$) mixtures ($n = 5,6,8,10,12$) have been obtained by the critical opalescence method using a laser scattering technique. All the systems show an upper critical solution temperature (UCST). In addition, the LLE curves have a rather horizontal top and their symmetry depends on the alkane size. The UCST increases almost linearly with n . For systems including a given alkane and phenol or phenylmethanol, the UCST is much higher than that of the corresponding mixtures with hexan-1-ol or heptan-1-ol. This reveals that dipolar interactions are stronger in solutions with aromatic alcohols. Preliminary DISQUAC interaction parameters for OH/aliphatic contacts in the investigated systems were obtained. It is remarkable that the coordinates of the critical points of (phenol or phenylmethanol + alkane) mixtures can be described using the same quasichemical interaction parameters for the OH/aliphatic and OH/aromatic contacts.

Keywords: liquid-liquid equilibria, phenylmethanol, alkane, DISQUAC

1. INTRODUCTION

Due to their complexity, the treatment of (alkanol + polar compound) mixtures is a severe test for any theoretical model. This can be ascribed to the alkanol self-association partially destroyed by the active molecules of the polar component, and to the new interactions between unlike molecules created upon mixing. Up to now, we have developed detailed studies of systems of the type (alkan-1-ol + polar compound), as those containing linear ketones,¹ linear organic carbonates,² linear esters,³ oxaalkanes,⁴⁻⁶ thiophene, tetrahydro-, 1,1-dioxide,⁷ or *N,N*-dialkylamides.^{8,9} Now, we extend these studies by replacing the alkan-1-ol by an aromatic alcohol. The aim of this series of articles is to investigate the effect of the aromatic ring on the thermodynamic properties of the mentioned solutions in order to gain insight into their structure and interactions. The dipolar moment in gas phase, μ , of aromatic alcohols, e.g., phenol, phenylmethanol, 2-methylphenol, is very similar to that of alkan-1-ols ($\approx 5.67 \cdot 10^{-30} \text{C}\cdot\text{m}$).¹⁰ In addition, they have slightly higher dielectric constants than alkan-1-ols with the same number of C atoms.¹¹⁻¹³ Thus, strong dipolar interactions may be expected in systems with aromatic alcohols. As a first step, we present here liquid-liquid equilibrium temperatures (T_{LLE}) as a function of the mole fraction (x_1) of phenylmethanol for systems containing heptane, octane, decane, dodecane or tetradecane as solvents. This type of data have been previously reported for the heptane solution.¹⁴

Aromatic alcohols show a large variety of relevant technical applications. Particularly, the properties of phenylmethanol have been the subject of considerable interest due to its versatility. For pharmaceutical aid, it is used as an antimicrobial agent;¹⁵ In addition, it is a good solvent for gelatine, cellulose acetate and shellac. Other applications are encountered in perfumery, veterinary science or in microscopy as embedding material.¹⁶ Phenylmethanol has been proposed as additive for synthetic fuels derived from the Fischer–Tropsch process for use in ground and air vehicles.¹⁷ It is also an additive that significantly retard the thermal degradation of jet fuels at high temperatures.¹⁷ Its applications as a dielectric solvent for the dielectrophoretic reconfiguration of nanowires are also remarkable.¹⁸

2. EXPERIMENTAL

2.1 Materials

Heptane (142-82-5, puriss p.a. ≥ 0.995); octane (11-65-9, puriss p.a. ≥ 0.99), decane (124-18-5, puriss p.a. ≥ 0.99), dodecane (112-40-3, puriss p.a. ≥ 0.98) and tetradecane (629-59-4, puriss p.a. ≥ 0.99) were from Fluka (purities expressed in mass fraction). Phenylmethanol (100-51-6, puriss p.a. ≥ 0.99) was from Aldrich. Prior to the measurements, the chemicals were stored over molecular sieves (Union Carbide Type 0.4 nm from Fluka). The densities, ρ , measured with an Anton-Paar densimeter DMA 602 (uncertainty $0.05 \text{ kg}\cdot\text{m}^{-3}$) at 298.15 K and

1
2
3 atmospheric pressure, were in good agreement with literature values as differences between
4 experimental and literature data are lower than 0.5% (Table 1). The water contents, determined
5 from the Karl-Fischer method, were (in mole fraction): 0.0003, 0.00012, 0.00011, 0.00003,
6 0.00015, and 0.00002 for phenylmethanol, heptane, octane, decane, dodecane, and tetradecane,
7 respectively.
8

9 10 2.2 Apparatus and Procedure

11 Mixtures were prepared by mass (weighing accuracy ± 0.01 kg), in small Pyrex tubes
12 (0.009 m i.d. and about 0.04 m length), immediately sealed by capping at atmospheric pressure
13 and room temperature. The relative atomic mass table of 1995 issued by IUPAC in 1996¹⁹ was
14 used for the conversion to molar quantities.
15

16 The liquid-liquid equilibrium curves for the investigated binary mixtures were
17 determined by the method of the critical opalescence. Details of the experimental technique are
18 given elsewhere.^{20,21} A Pt-1000 resistance was used for the measurement of the T_{LLE} values.
19 The Pt-1000 resistance was calibrated, according to the ITS-90 scale of temperature, against the
20 triple point of the water and the fusion point of Ga. The precision of the equilibrium temperature
21 measurements is ± 0.001 K, with an estimated uncertainty of ± 0.05 K. The mentioned
22 temperatures were reproducible to ± 0.02 K for those values close to the UCST. For the
23 equilibrium composition measurements, the uncertainty of the mole fraction is better than
24 0.0005. As a matter of fact, the precision of the weighing technique is 0.0001 in mole fraction,
25 but, because the more volatile component is partially evaporated to the free volume of the
26 ampoule ($\approx 1.17 \cdot 10^{-6}$ m³), this value is slightly reduced.
27
28
29
30
31
32
33
34
35
36

37 3. RESULTS

38 Table 2 lists the directly measured T_{LLE} values obtained in this work vs. x_1 for
39 (phenylmethanol + heptane, + octane, + decane, + dodecane, or + tetradecane) mixtures (see
40 Figs. 1-3).
41

42 All the systems show an UCST. Note that the LLE curves have a flat maximum, and
43 that their symmetry depends on the alkane size (Fig. 1).
44

45 Table 3 lists the composition (x_{1c}) and temperature (T_c) of the critical points obtained
46 by reducing the experimental measurements using the equation^{22,23}
47

$$48 \quad T / K = T_c / K + k |y - y_c|^m \quad (1)$$

49 being
50

$$51 \quad y = \frac{\alpha x_1}{1 + x_1(\alpha - 1)} \quad (2)$$

52
53
54
55
56
57
58
59
60

$$y_c = \frac{\alpha x_{1c}}{1 + x_{1c}(\alpha - 1)} \quad (3)$$

In equations (1) to (3), m , k , α , T_c and x_{1c} are the parameters to be fitted against the experimental data. For $\alpha = 1$, equation (1) is similar to:²⁴⁻²⁶

$$\Delta\lambda = B\tau^\beta \quad (4)$$

where, $\Delta\lambda_1 = \lambda_1' - \lambda_2''$ is the so-called order parameter, which is any density variable in the conjugate phase (in the present case, $\lambda_1 = x_1$), $\tau (= T_c - T)/T_c$ is the reduced temperature and β denotes the critical exponent connected to $\Delta\lambda_1$. The β value depends on the theory applied to its determination.²⁴⁻²⁶ More details can be encountered elsewhere.²⁷

The adjustment of the parameters involved in equation (1) was performed on the basis of the Marquardt algorithm²⁸ with all the points weighted equally. Values of the fitted m , k , α , T_c , x_{1c} parameters are collected in Table 3, which also contains the standard deviations for temperatures calculated according to:

$$(\sigma(T_{LLE})/K) = \left[\sum (T_{LLEi}^{\text{exp}} - T_{LLEi}^{\text{cal}})^2 / (N - n) \right]^{1/2} \quad (5)$$

Here, N and n are, respectively, the number of data points and the number of fitted parameters. Equation (1) fits well the experimental measurements. Our coordinates for the critical point of the heptane solution are close to literature values ($x_{1c} = 0.4866$; $T_c = 324.65$ K).¹⁴

4. DISCUSSION

The UCST of the studied systems increases with the number of C atoms in the chain length of the alkane (Fig. 2, Table 3). The same behaviour is observed in mixtures formed by alkane with, azapan-2-one,²⁰ linear alkanone,²⁹ linear organic carbonate,³⁰ acetic anhydride,³¹ alkoxyethanol,^{27,32,33} polyether,^{34,35} or phenol.³⁶⁻⁴⁰ Fig. 3 shows that the LLE curves are progressively shifted to higher x_1 values when the number of C atoms of the alkane increases. A similar trend is encountered for many other mixtures.^{27,31-40}

It is remarkable that mixtures involving phenylmethanol or phenol and a given alkane are characterized by close UCST values. Thus, for heptane systems, UCST(phenol) = 326.05 K³⁶ or 327.3;³⁸ UCST(phenyl methanol) = 323.67 K, for mixtures with octane, UCST(phenol)³⁹ = 329.54 K; UCST(phenyl methanol) = 326.85 K; and for solutions containing decane,

UCST(phenol)³⁹ = 336.50 K; UCST(phenyl methanol) = 335.71 K. This reveals that in such mixtures interactions between like molecules are rather similar. In contrast, UCSTs of systems involving phenol or phenylmethanol are much higher than those of mixtures with hexan-1-ol or heptan-1-ol, as these alcohols are miscible with alkanes at 298.15 K at any composition.^{41,42} This indicates that alkanol-alkanol interactions are enhanced by the aromatic ring of phenol or phenylmethanol. A similar behaviour is observed, e.g., for (1-hexanamine or aniline + a given alkane) systems. Thus, at equimolar composition and 298.15 K, for (1-hexanamine + heptane),⁴³ the molar excess enthalpy is 1064 J•mol⁻¹, while UCST(aniline + heptane)³⁷ = 343.11 K.

It is known that the impact of polarity on bulk properties may be characterized by using the effective dipole moment, $\bar{\mu}$.^{24,41,45} This magnitude is defined by:

$$\bar{\mu} = \left[\frac{\mu^2 N_A}{4\pi\epsilon_0 V k_B T} \right]^{1/2} \quad (6)$$

where N_A is the Avogadro's number, k_B , the constant Boltzmann; V , the molar volume, and ϵ_0 the permittivity of the vacuum. For a given homologous series (e.g. alkan-1-ols) at temperature T , application of equation (6) leads to a much higher variation with compound size of $\bar{\mu}$ compared to that observed for μ .⁴⁵ Thus at 298.15 K, $\bar{\mu}$ (methanol) = 1.023 > $\bar{\mu}$ (ethanol) = 0.852 > $\bar{\mu}$ (butan-1-ol) = 0.664 > $\bar{\mu}$ (hexan-1-ol) = 0.580 > $\bar{\mu}$ (heptan-1-ol) = 0.547 > $\bar{\mu}$ (decan-1-ol) = 0.443. For phenylmethanol, $\bar{\mu}$ = 0.639 (value calculated using the density listed in Table 1 and $\mu = 5.67 \cdot 10^{-30} \text{ C}\cdot\text{m}^{10}$). For phenol, $\bar{\mu}$ = 0.647 (value obtained at 318.15 K using density reported in the literature¹¹ and $\mu = 5.50 \cdot 10^{-30} \text{ C}\cdot\text{m}^{10}$). The difference for $\bar{\mu}$ values of these aromatic alcohols with those of hexan-1-ol and heptan-1-ol may explain the lower solubility of the former in alkanes.

Finally, we have explored the possibility of representing the (x_{1c}, T_c) values of the phenylmethanol solutions considered here by means of the group contribution model DISQUAC^{44,45} on the basis of our previous study on phenol systems.⁴⁵ Details on DISQUAC have been reported elsewhere.^{44,45} The geometrical parameters of the groups referred to in this work (see below) are available in the literature.⁴⁶ The variation with the temperature of the interaction parameters is expressed in terms of the DIS (dispersive) and QUAC (quasichemical) interchange coefficients,^{44,45} $C_{st,l}^{\text{DIS}}; C_{st,l}^{\text{QUAC}}$ where $s \neq t$ are two contact surfaces present in the mixture and $l = 1$ (Gibbs energy); $l = 2$ (enthalpy); $l = 3$ (heat capacity). The investigated mixtures are built by three surfaces: type a, aliphatic (CH₃, CH₂, in alkanes, or phenylmethanol); type b, aromatic (C₆H₅ in phenylmethanol) and type h, hydroxyl (OH in phenylmethanol). Thus,

(phenylmethanol + alkane) mixtures are built by three types of contact: (a,b); (a,h) and (b,h). The interaction parameters are entirely dispersive for the (a,b) contact and are known from the study of (alkyl benzene + alkane) mixtures.⁴⁷ The (a,h) and (b,h) contacts are characterized by DIS and QUAC interchange coefficients. The values for $C_{bh,1}^{DIS}$, $C_{sh,1}^{DIS}$ ($s = a,b; l = 2,3$), and for $C_{sh,l}^{QUAC}$ ($s = a,b$ and $l = 1,2,3$) are assumed to be the same as in phenol mixtures. Therefore, only the coefficients $C_{ah,1}^{DIS}$ have been fitted in order to represent the data. Values of the final interchange coefficients are given in Table 4. DISQUAC results for the coordinates of the critical points are listed in Table 3. It is remarkable that they are correctly represented (Table 3), as differences between calculated and experimental UCST do not exceed 5.5 K, and for x_{1c} such differences are 0.1 or lower. Discrepancies between experimental and calculated results, as well as the variation of the $C_{ah,1}^{DIS}$ ($l = 1,3$) coefficients with the size of the alkane can be explained as follows. DISQUAC is a mean field theory and, in this type of theories, excess functions are assumed to be analytical close to the critical points, in such way that calculations are developed under this basic (and erroneous) assumption. In fact, thermodynamic properties, in the proximity of the critical point, follow scaling laws with universal critical exponents and universal scaling functions.²⁴ According to the critical exponents theory,²⁴ DISQUAC provides LLE curves which are: (i) more rounded than the experimental ones at temperatures near to the UCST; (ii) too high at the UCST and too low at the LCST²⁴ (lower critical solution temperature). Thus, the $C_{ah,1}^{DIS}$ ($l = 1,3$) coefficients must be kept between certain limits in order to provide not very high calculated values of UCSTs. The same occurs in other applications as those on systems including thiophene, tetrahydro-, 1,1-dioxide,⁷ alkoxyethanol,⁴⁹ amide,⁸ or pyridine⁵⁰ and alkane. The theoretical results obtained here suggest that systems with phenol or phenylmethanol can be described using the same QUAC interaction parameters for the OH/aliphatic and OH/aaromatic contacts. A similar behaviour is encountered for (aniline⁴⁹ or 2-methylaniline²¹ + alkane) solutions.

5. CONCLUSIONS

Liquid-liquid equilibrium curves were determined for mixtures of phenyl methanol with heptane, octane, decane, dodecane and tetradecane. For systems with a given alkane and phenol or phenylmethanol, the UCST is much higher than that of the corresponding mixtures with hexan-1-ol or heptan-1-ol, which reveals that dipolar interactions are stronger in solutions with aromatic alcohols. Preliminary DISQUAC interaction parameters for the OH/aliphatic contacts in the investigated systems were obtained. The $C_{sh,l}^{QUAC}$ ($s = a,b; l = 1,2,3$) coefficients are the same as in phenol + alkane mixtures.

6. LITERATURE CITED

- (1) González, J.A. Thermodynamics of mixtures containing the CO and OH groups. I. Estimation of the DISQUAC interchange coefficients for 1-alkanols + *n*-alkanones systems. *Can. J. Chem.*, 1997, 75, 1412-1423.
- (2) González, J.A.; Szurgocinska, M.; Domanska, U. Thermodynamics of mixtures containing organic carbonates. Part XIII. Solid-liquid equilibria of long-chain 1-alkanol + dimethyl or diethyl carbonate systems. DISQUAC and ERAS analysis of the hydroxyl/carbonate interactions. *Fluid Phase Equilib.*, 2002, 200, 349-374.
- (3) González, J.A.; Mozo, I.; García de la Fuente, I.; Cobos, J.C. Thermodynamics of 1-alkanol + linear alkanoate mixtures. *Phys. Chem. Liq.*, 2005, 43, 175-194.
- (4) Cobos, J.C.; García, I.; González, J.A.; Casanova, C. Excess enthalpies of (3,6,9-trioxaundecane + an *n*-alkan-1-ol) at 298.15 K. *J. Chem. Thermodyn.*, 1990, 22, 383-386
- (5) González, J.A.; Mozo, I.; García de la Fuente, I.; Cobos, J.C.; Riesco, N. Thermodynamics of (1-alkanol + linear monoether) systems. *J. Chem. Thermodyn.*, 2008, 40, 1495-1598.
- (6) González, J.A.; Mozo, I.; García de la Fuente, I.; Cobos, J.C.; Durov, V.A. Thermodynamics of 1-alkanol + cyclic ether mixtures. *Fluid Phase Equilib.*, 2006, 245 168-184.
- (7) González, J.A.; Domanska, U. Thermodynamics of mixtures containing a very strongly polar compound. Part I. Experimental phase equilibria (solid-liquid and liquid-liquid) for sulfolane + alkan-1-ols systems. Analysis of some mixtures including sulfolane in terms of DISQUAC. *Phys. Chem. Chem. Phys.*, 2001, 3, 1034-1042.
- (8) González, J.A.; Cobos, J.C.; García de la Fuente, I. Thermodynamics of liquid mixtures containing a very strongly polar compound. Part 6. DISQUAC characterization of *N,N*-dialkylamides. *Fluid Phase Equilibria*, 2004, 224, 169-183.
- (9) González, J.A.; Domanska, U.; Lachwa, J. Thermodynamics of mixtures containing a very strongly polar compound. Part 3. DISQUAC characterization of NMP + organic solvent mixtures. *Can. J. Chem.*, 2003, 81, 1-11.
- (10) McClellan, A.L. *Tables of Experimental Dipole Moments*, Rahrha Enterprises, El Cerrito, CA, 1974.
- (11) Riddick, J. A.; Bunger, W. B.; Sakano, T. K., *Organic Solvents, Techniques of Chemistry Vol 2*, 1986.
- (12) Iglesias, T.P.; Legido, J.L.; Pereira, S.M.; Comings, B.; Paz Andrade, M.I. Relative permittivities and refractive indices on mixing for (*n*-hexane + 1-pentanol, or 1-hexanol, or 1-heptanol) at $T = 298.15$ K. *J. Chem. Thermodyn.*, 2000, 32, 923-930.

- 1
2
3 (13) Piritä-Honkanen, P.; Ruostesuo, R. Thermodynamic and spectroscopic properties of 2-
4 pyrrolidone. Part 5. Dielectric properties of 2-pyrrolidone + benzyl alcohol, + 1,4-dioxane
5 and + hexamethylphosphortriamide at 303.15 and 323.15 K. *Thermochim. Acta*, **1991**,
6 *184*, 65-71.
7
8
9 (14) Hughes, M. A.; Haoran, Y. Liquid-liquid equilibria for separation of toluene from heptane
10 by benzyl alcohol tri(ethylene glycol) mixtures. *J. Chem. Eng. Data* **1990**, *35* 467-471.
11
12 (15) Sweetman, S., *Martindale: The complete drug reference*. 37th ed.; Pharmaceutical Press:
13 London, **2011**.
14
15 (16) *The Merck Index*. 14th ed.; Merck Publishing: **2006**.
16
17 (17) Link, D. D.; Gormley, R. J.; Baltrus, J. P.; Anderson, R. R.; Zandhuis, P. H. Potential
18 additives to promote seal swell in synthetic fuels and their effect on thermal stability.
19 *Energy & Fuels* **2008**, *22*, 1115-1120.
20
21 (18) Wissner-Gross, A. Dielectrophoretic reconfiguration of nanowire interconnects.
22 *Nanotechnology*, **2006**, *17*, 4986-4990.
23
24 (19) IUPAC: Atomic weights of the elements 1995 (IUPAC technical report). *Pure Appl.*
25 *Chem.*, **1996**, *68*, 2339-2359.
26
27 (20) Alonso, V.; Alonso, I.; Mozo, I.; González, J. A.; De La Fuente, I. G.; Cobos, J. C.
28 Thermodynamics of mixtures containing a strongly polar compound. 9. liquid-liquid
29 equilibria for ϵ -caprolactam + selected alkanes. *J. Chem. Eng. Data* **2010**, *55*, 2263-2266.
30
31 (21) Fernández Regúlez, M; Mozo, I.; González, J.A.; García de la Fuente, I.; Cobos, J.C.
32 Thermodynamics of organic mixtures containing amines: VI. Liquid-liquid equilibria
33 for mixtures of *o*-toluidine + selected alkanes. *Fluid Phase Equilib.*, **2007**, *259*,39-44.
34
35 (22) Ott, J. B.; Holscher, I. F.; Schneider, G. M., (Liquid-Liquid) phase equilibria in (methanol
36 + heptane) and (methanol + octane) at pressures from 0.1 to 150 MPa. *J. Chem.*
37 *Thermodyn.* **1986**, *18*, 815-826.
38
39 (23) Haarhaus, U.; Schneider, G. M., (Liquid + liquid) phase equilibria in (methanol + butane)
40 and (methanol + pentane) at pressures from 0.1 to 140 MPa. *J. Chem. Thermodyn.* **1988**,
41 *20*, 1121-1129.
42
43 (24) Rowlinson, J. S.; Swinton, F. L., *Liquids and Liquid Mixtures*, **1982**.
44
45 (25) Stanley, H. E., *Introduction to Phase Transitions and Critical Phenomena*, **1971**.
46
47 (26) Novak, J. P.; Matous, J.; Pick, J., *Liquid-Liquid Equilibria*, **1987**.
48
49 (27) Rubio, M. A.; González, J. A.; García De La Fuente, I.; Cobos, J. C. Thermodynamic
50 properties of *n*-alkoxyethanols + organic solvents mixtures VIII. Liquid-liquid equilibria of
51 systems containing 2-methoxyethanol and alkanes (C₆H₁₂ and CH₃-(CH₂)_u-CH₃, u =
52 3,4,6,8). *Fluid Phase Equilib.* **1998**, *143*, 111-123.
53
54 (28) Bevington, P. R., *Data Reduction and Error Analysis for the Physical Sciences*, **1969**.
55
56
57
58
59
60

- 1
2
3 (29) Messow, U.; Doyé, U.; Kuntzsch, S. Liquid-liquid equilibrium, excess enthalpies and
4 excess volumes of some alkanone (C3, C4) + normal alkane (C10, C12, C14, C16)
5 mixtures. *Int. DATA Ser., Sel. Data Mixtures, Ser. A* **1979**, *1*, 151-158.
6
7 (30) Gonzalez, J. A.; Garcia, I.; Cobos, J. C.; Casanova, C. Thermodynamics of binary mixtures
8 containing organic carbonates. 4. Liquid-liquid equilibria of dimethyl carbonate + selected
9 *n*-alkanes. *J. Chem. Eng. Data* **1991**, *36*, 162-164.
10
11 (31) Aboy, M.; Villa, S.; Riesco, N.; González, J. A.; De La Fuente, I. G.; Cobos, J. C. Liquid-
12 liquid equilibria for acetic anhydride + selected alkanes. *J. Chem. Eng. Data*, **2002**, *47*,
13 950-953.
14
15 (32) Martínez, R.; Gonzalez, J. A.; Garcia De La Fuente, I.; Cobos, J. C. Thermodynamic
16 properties of *n*-alkoxyethanols+organic solvent mixtures. XIV. Liquid-liquid equilibria of
17 systems containing 2-(2-ethoxyethoxy)ethanol and selected alkanes. *J. Chem. Eng. Data*,
18 **2000**, *45*, 1036-1039.
19
20 (33) Rubio, M. A.; González, J. A.; García De La Fuente, I.; Cobos, J. C. Thermodynamic
21 properties of *n*-alkoxyethanols + organic solvent mixtures. IX. Liquid-liquid equilibria of
22 systems containing 2-methoxyethanol or 2-ethoxyethanol and selected *n*-alkanes. *J. Chem.*
23 *Eng. Data*, **1998**, *43*, 811-814.
24
25 (34) Mozo, I.; González, J. A.; De La Fuente, I. G.; Cobos, J. C. Thermodynamics of mixtures
26 containing ethers. Part III. liquid-liquid equilibria for 2,5,8,11-tetraoxadodecane or
27 2,5,8,11,14-pentaoxapentadecane + selected *n*-alkanes. *J. Chem. Eng. Data*, **2004**, *49*,
28 1091-1094.
29
30 (35) Treszczanowicz, T.; Cieślak, D. (Liquid + liquid) equilibria in (a dimethyl ether of a
31 polyethylene glycol + an *n*-alkane). *J. Chem. Thermodyn.*, **1993**, *25*, 661-665.
32
33 (36) Vondracek, R. On binary systems of phenol with some hydrocarbons. *Collect. Czech.*
34 *Chem. Commun.*, **1937**, *9*, 168-175.
35
36 (37) Matsuda, H.; Ochi, K.; Kojima, K.. Determination and correlation of LLE and SLE data for
37 methanol + heptane, aniline + heptane and phenol + hexane system. *J. Chem. Eng. Data*,
38 **2003**, *48*, 184-189.
39
40 (38) Matsuda, H.; Fujita, M.; Ochi, K. Measurement and correlation for high-viscosity binary
41 systems: aniline + methylcyclohexane, phenol + heptane, phenol + octane, and glycerol +
42 1-pentanol. *J. Chem. Eng. Data*, **2003**, *48*, 1076-1080.
43
44 (39) Gmehling, J. Phase equilibria in binary systems formed by phenol with benzene, *n*-octane,
45 and *n*-decane. *J. Chem. Eng. Data*, **1982**, *27*, 371-373.
46
47 (40) Ksiazczak, A.; Kosinski, J.J. Liquid-liquid equilibrium in binary polar aromatic +
48 hydrocarbon systems. *Fluid Phase Equilib.*, **1991**, *59*, 291-308.
49
50 (41) Christensen, J.J.; Izatt, R.M.; Stitt, B.D.; Hanks, R.W. The excess enthalpies of seven *n*-
51 decane + alcohol mixtures at 298.15 K. *J. Chem. Thermodyn.*, **1979**, *11*, 261-266.
52
53
54
55
56
57
58
59
60

- 1
2
3 (42) Amigo, A.; Legido, J.L.; Bravo, R.; Paz Andrade, M.I. Excess molar enthalpies of
4 (heptan-1-ol + an *n*-alkane) at 298.15 and 308.15 K. *J. Chem. Thermodyn.*, **1989**, *21*,
5 1207-1211.
6
7 (43) Pfestorf, R.; Kuchenbecker, D.; Quitzsch, K. Kalorische Exzesscharakteristik flüssiger
8 Nichtelektrolytsysteme. IV. Experimentelle Bestimmung und Vorausberechnung
9 mittlerer molarer Mischungsenthalpien von *n*-Alkylamin/*n*-Alkan-Systemen. *Z. Phys.*
10 *Chem. (Leipzig)*, **1982**, *263*, 233-240.
11
12 (44) González, J.A.; García de la Fuente, I.; Cobos, J.C. Proximity effects and cyclization in
13 oxaalkanes + CCl₄ mixtures. DISQUAC characterization of the Cl-O interactions.
14 Comparison with Dortmund UNIFAC results. *Fluid Phase Equilib.*, **1999**, *154*, 11-31.
15
16 (45) González, J.A.; García de la Fuente, I.; Cobos, J.C. Thermodynamics of mixtures with
17 strongly negative deviations from the Raoult's law. Part 4. Application of the
18 DISQUAC model to mixtures of 1-alkanols with primary or secondary linear amines.
19 Comparison with Dortmund UNIFAC and ERAS results. *Fluid Phase Equilib.*, **2000**,
20 *168*, 31-58.
21
22 (46) González, J.A.; García de la Fuente, I.; Cobos, J.C. Application of a purely physical
23 model (DISQUAC) to binary mixtures of phenol and organic solvents. *Ber. Bunsenges.*
24 *Phys. Chem.*, **1996**, *100*, 1746-1751.
25
26 (47) Cannas, A.; Marongiu, B.; Porcedda, S. Thermodynamic properties of *n*-alkylbenzene +
27 *n*-alkane or cyclohexane mixtures. Comparison with DISQUAC predictions.
28 *Thermochim. Acta*, **1998**, *311*, 1-19.
29
30 (48) González, J.A.; Cobos, J.C.; Carmona, F.J.; García de la Fuente, I.; Bhethanabotla,
31 V.R.; Campbell, S. W. Thermodynamics of systems containing alkoxyethanols. Part
32 XV. DISQUAC characterization of systems of alkoxyethanols with *n*-alkanes or
33 cyclohexane. *Phys. Chem. Chem. Phys.*, **2001**, *3*, 2856-2865.
34
35 (49) González, J.A.; Mozo, I.; García de la Fuente, I.; Cobos, J.C. Thermodynamics of
36 organic mixtures containing amines. V. Systems with pyridines. *Thermochim. Acta*,
37 **2006**, *441*, 53-68.
38
39 (50) González, J.A.; Mozo, I.; García de la Fuente, I.; Cobos, J.C. Thermodynamics of
40 organic mixtures containing amines. IV. Systems with aniline. *Can. J. Chem.*, **2005**, *83*,
41 1812-1825
42
43 (51) Zhu, S.; Shen, S.; Benson, G. C.; Lu, B. C. Y. Excess enthalpies of (methyl 1,1-
44 dimethylpropyl ether + an *n*-alkane) at the temperature 298.15 K. *J. Chem. Thermodyn.*,
45 **1994**, *26*, 35-39.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ACKNOWLEDGEMENTS

The authors gratefully acknowledge the financial support received from the Ministerio de Educación y Ciencia, under the Project FIS2007-61833.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

TABLE 1**Comparison of Experimental Densities, ρ , with Literature for Pure Liquids at $T = 298.15$ K.**

Compound	$\rho/\text{kg}\cdot\text{m}^{-3}$	
	Exp.	Lit.
Phenylmethanol	1043.53	1041.30 ¹¹
Heptane	679.71	679.46 ¹¹
Octane	698.76	698.62 ¹¹
Decane	726.43	726.35 ¹¹
Dodecane	741.54	745.18 ¹¹
Tetradecane	759.27	759.29 ⁵¹

^a Density uncertainty: 0.05 kg·m⁻³

TABLE 2

Experimental Liquid-liquid Equilibrium Temperatures (T_{LLE}) for (Phenylmethanol(1) + Alkane(2)) Mixtures^a.

X_1	T_{LLE}/K	x_1	T_{LLE}/K
Phenylmethanol(1) + heptane(2)			
0.2897	321.00	0.4744	323.74
0.2968	321.39	0.4875	323.71
0.3159	321.99	0.4953	323.72
0.3319	322.37	0.5019	323.68
0.3536	322.88	0.5223	323.63
0.3537	322.89	0.5384	323.54
0.3832	323.24	0.5520	323.48
0.3923	323.37	0.5657	323.36
0.3957	323.40	0.5702	323.35
0.3984	323.41	0.5822	323.25
0.4059	323.52	0.5988	322.89
0.4336	323.63	0.6036	322.89
0.4384	323.73	0.6214	322.50
0.4478	323.64	0.6297	322.13
0.4638	323.76	0.6685	319.74
0.4683	323.70		
Phenylmethanol(1) + octane(2)			
0.2715	321.80	0.5219	326.79
0.2853	322.47	0.5266	326.86
0.3022	323.22	0.5423	326.74
0.3136	323.74	0.5555	326.73
0.3259	324.23	0.5755	326.69
0.3312	324.43	0.6016	326.52
0.3435	324.87	0.6133	326.42
0.3526	325.19	0.6163	326.34
0.3698	325.54	0.6395	325.99
0.3763	325.69	0.6499	325.82
0.3822	325.83	0.6647	325.38
0.3996	326.19	0.6752	324.99
0.4096	326.24	0.6896	324.34
0.4223	326.47	0.7044	323.68
0.4416	326.58	0.7046	323.62

TABLE 2 (continued)

0.4636	326.71	0.7152	323.01
0.4914	326.82	0.7237	322.58
0.5198	326.86		
Phenylmethanol 1) + decane(2)			
0.2736	326.26	0.5898	335.81
0.3108	329.37	0.6033	335.73
0.3193	329.85	0.6055	335.68
0.3436	331.36	0.6271	335.67
0.3622	332.20	0.6357	335.62
0.3981	333.57	0.6668	335.57
0.4349	334.62	0.6971	335.14
0.4534	334.89	0.7051	334.88
0.4738	335.38	0.7095	334.84
0.4750	335.23	0.7229	334.45
0.5094	335.53	0.7255	334.32
0.5290	335.66	0.7404	333.80
0.5379	335.64	0.7441	333.57
0.5578	335.74	0.7495	333.23
0.5724	335.75	0.7888	330.71
0.5779	335.79		
Phenylmethanol(1) + dodecane(2)			
0.3605	338.59	0.6395	345.09
0.3743	339.40	0.6544	345.04
0.3936	340.22	0.6780	345.05
0.4044	340.80	0.6904	344.99
0.4182	341.48	0.7086	344.91
0.4440	342.47	0.7229	344.77
0.4617	343.02	0.7387	344.57
0.4883	343.79	0.7442	344.48
0.4911	343.77	0.7576	344.21
0.5114	344.19	0.7656	343.99
0.5255	344.41	0.7786	343.29
0.5448	344.65	0.7865	343.12
0.5639	344.83	0.7980	342.51

TABLE 2 (continued)

0.5833	344.90	0.8151	341.29
0.6064	344.99	0.8420	338.47
0.6207	345.00		
	Phenylmethanol(1) + tetradecane(2)		
0.4502	349.47	0.6212	353.83
0.4544	349.71	0.6344	353.90
0.4702	350.39	0.6560	353.93
0.4863	351.06	0.6803	354.00
0.4965	351.31	0.7000	353.99
0.5091	351.861	0.7173	353.94
0.5158	351.99	0.7386	353.89
0.5170	352.09	0.7517	353.89
0.5265	352.31	0.7778	353.55
0.5485	352.85	0.7993	353.09
0.5631	353.08	0.8041	352.82
0.5695	353.19	0.8108	352.62
0.5832	353.47	0.8232	352.02
0.5966	353.60		

^a uncertainties, u are; $u(x_1) = 0.0005$; $u(T_{\text{LLE}}) = 0.05$ K

TABLE 3

Coefficients in Eq. (1) for the Fitting of the (x_1, T_{LLE}) Pairs Given in Table 2 for
(Phenylmethanol(1) + Alkane(2)) Mixtures; σ is the Standard Deviation Defined by Eq 5.

N^a	m	K	α	T_c/K	x_{1c}	σ/K
Phenylmethanol(1) + heptane(2)						
31	3.305	-1065	0.564	323.67 (327.0) ^b	0.487 (0.377) ^b	0.07
Phenylmethanol(1) + octane(2)						
35	2.740	-283	0.768	326.85 (331.3) ^b	0.518 (0.421) ^b	0.07
Phenylmethanol(1) + decane(2)						
31	3.248	-583	0.652	335.71 (341.1) ^b	0.583 (0.507) ^b	0.07
Phenylmethanol(1) + dodecane(2)						
31	3.190	-528	0.460	345.03 (349.9) ^b	0.647 (0.589) ^b	0.04
Phenylmethanol(1) + tetradecane(2)						
27	3.190	-501	0.39128	353.97 (358.2) ^b	0.694 (0.647) ^b	0.015

^a number of experimental data points; ^bDISQUAC value calculated with the interchange coefficients listed in Table 4.

TABLE 4

Dispersive (DIS) and Quasichemical (QUAC) Interchange Coefficients, $C_{st,l}^{DIS}$ and $C_{st,l}^{QUAC}$, for (s,t) Contacts in (Phenylmethanol + Alkane) Mixtures ($l = 1$, Gibbs Energy; $l = 2$, Enthalpy; $l = 3$, Heat Capacity).

Contact ^a (s,t)	$C_{st,1}^{DIS}$	$C_{st,2}^{DIS}$	$C_{st,3}^{DIS}$	$C_{st,1}^{QUAC}$	$C_{st,2}^{QUAC}$	$C_{st,3}^{QUAC}$
(b,h)	4.80	-2.65		5.70	13	
(a,h) $n \leq 7$ ^b	3.42	0.50	-5	11.25	16	12
(a,h) $n = 8$ ^b	3.32	0.50	-5	11.25	16	12
(a,h) $n > 8$ ^b	3.32	0.50	5	11.25	16	30

^aa, aliphatic in alkane, or phenylmethanol; b, C₆H₅ in phenylmethanol; h, OH, in phenylmethanol; ^b n is the number of C atoms in the alkane




Figure 1

CAPTION TO FIG. 1

LLE of (phenylmethanol(1) + alkane(2)) mixtures. Points, experimental results: (■), heptane; (●), octane; (▲), decane; (□), dodecane; (*), tetradecane. Solid lines, results from the fitting equation (1).




Figure 2

CAPTION TO FIG. 2

Upper critical solution temperatures T_c , vs. n , the number of carbon atoms of the alkane, for (phenylmethanol + alkane) mixtures.





Figure 3

CAPTION TO FIG. 3

Upper critical solution composition, x_{1c} , vs. n , the number of carbon atoms of the alkane, for (phenylmethanol + alkane) mixtures.

1
2
3
4
5
6
7
8
9 **For Table of Contents use only**



25 **THERMODYNAMICS OF MIXTURES CONTAINING AROMATIC ALCOHOLS. 1.**
26 **LIQUID-LIQUID EQUILIBRIA FOR (PHENYLMETHANOL + ALKANE) SYSTEMS**

27
28
29
30 JUAN ANTONIO GONZÁLEZ, CRISTINA ALONSO-TRISTÁN, ISAÍAS GARCÍA DE LA
31 FUENTE AND JOSÉ CARLOS COBOS
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60