

Excess enthalpies of binary and ternary mixtures containing dibutyl ether (DBE), cyclohexane and 1-butanol at 298.15 K

Journal:	<i>Journal of Chemical & Engineering Data</i>
Manuscript ID:	je-2008-00751m.R1
Manuscript Type:	Article
Date Submitted by the Author:	13-Jan-2009
Complete List of Authors:	Aguilar, Fernando; Universidad de Burgos, Ingeniería Electromecánica Alaoui, Fatima; Universidad de Burgos, Ingeniería Electromecánica Alonso-Tristán, Cristina; Universidad de Burgos, Ingeniería Electromecánica SEGOVIA, JOSÉ; UNIVERSITY OF VALLADOLID, INGENIERÍA ENERGÉTICA Y FLUIDOMECAÁNICA Villamanan, Miguel; Universidad de Valladolid, Depto. de Ingeniería Energetica y Fluidomecánica Montero, Eduardo; Universidad de Burgos, Ingeniería Electromecánica



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Excess enthalpies of binary and ternary mixtures containing dibutyl ether (DBE), cyclohexane and 1- butanol at 298.15 K¹.

Fernando Aguilar^(a), Fatima E.M. Alaoui^(a), Cristina Alonso-Tristán^(a), José J. Segovia^(b), Miguel A. Villamañán^(b) and Eduardo A. Montero^{(a),}*

Departamento de Ingeniería Electromecánica, Escuela Politécnica Superior, Universidad de Burgos, E-09006 Burgos, Spain, Grupo de Termodinámica y Calibración TERMOCAL, E.T.S. de Ingenieros Industriales, Universidad de Valladolid, E-47071 Valladolid, Spain.

RECEIVED DATE (to be automatically inserted after your manuscript is accepted if required according to the journal that you are submitting your paper to)

*Corresponding author. E-mail: emontero@ubu.es. Tel. +34 947 258 916. Fax: +34 947 259 088

^(a) Universidad de Burgos

^(b) Universidad de Valladolid

¹ This paper is part of the Doctoral Thesis of F. Aguilar.

Experimental excess enthalpies of the ternary system dibutyl ether (DBE) + cyclohexane + 1-butanol and the corresponding binary systems at 298.15 K are reported. A quasi-isothermal flow calorimeter has been built and tested to make the measurements. All the binary and the ternary systems show endothermic character. The experimental data have been fitted using a polynomial equation for the

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

binary and ternary systems. The values of the standard deviation indicate good agreement between the experimental results and these calculated from the equation.

Introduction

The use of oxygenated compounds as gasoline-blending agents has been proposed to reduce emissions of new reformulated gasoline. Ether + alkane + alcohol mixtures are of interest as model mixtures for gasoline in which the alcohol and the ether act as non-polluting, high octane number blending agents. From this point of view the study of the ternary mixture dibutyl ether + cyclohexane + 1-butanol is very interesting. Cyclohexane is an usual compound of gasoline; dibutyl ether (DBE) is used as blending agent in reformulated gasoline and 1-butanol is a basic component in the synthesis of the ether, and therefore is always contained as an impurity. Experimental excess enthalpies of the ternary system DBE + cyclohexane + 1-butanol and the corresponding binary systems at 298.15 K are reported in this work. Excess enthalpies have been measured with a new quasi-isothermal flow calorimeter recently built at the University of Burgos. The experimental data have been fitted using a Redlich-Kister¹ equation and NRTL² and UNIQUAC³ models. The values of the standard deviation indicate the agreement between the experimental results and these calculated from the equations.

Experimental Section

Materials. All the chemicals used here were purchased from Fluka Chemie AG and were of the highest purity available, chromatography quality reagents (of the series puriss p.a.) with a stated purity >99.5 %. All reagents were checked by gas chromatography, and the values of purity obtained were >99.6 % for DBE, >99.9 % for cyclohexane and >99.8 % for 1-butanol. The water content for 1-butanol was checked to be less than 0.009%.

Apparatus and Procedure. Excess enthalpies have been measured with a new quasi-isothermal flow calorimeter recently built at the University of Burgos, based on the previous experience of the research group.^{4,5} Figure 1 shows schematically the quasi-isothermal calorimeter. Two precision positive displacement pumps (Agilent, model 1100) deliver the liquids through stainless-steel tubing at

programmable constant flow rates into the mixing coil sitting in the flow cell, which is included in the measure unit, following the scheme shown in Figure 2. The flow cell has been built in the TERMOCAL laboratory at the University of Valladolid. In case of binary systems, the two liquids are the pure components and, in case of ternary systems, one of them is a pure component and the other the corresponding binary mixture. The flow cell is immersed in a water bath, (Hart Scientific, model 6020E) thermostatted at the temperature the mixing process is taking place. The stability of the temperature is better than ± 0.01 K. The bath temperature is measured by its calibrated standard PRT-100, resolving 10 mK in the reading of temperature and estimating an overall uncertainty of ± 30 mK.

Isothermal calorimetry is based on measuring the energy required to maintain the mixing vessel at a constant temperature. To achieve this condition, a Peltier cooler removes at a constant rate energy from the flow cell and a control-heater compensates this energy and additionally the energy liberated (exothermic) or absorbed (endothermic) by the mixing process and maintains the temperature of the flow cell constant. The variation of temperature is detected by a control sensor, a NTC thermistor. The resistance is measured by a 4-wire resistance bridge with a micro-ohm meter from Hewlett-Packard, model HP-34420A. The Peltier cooler, the control-heater and the calibration heater are connected, each one, to its respectively DC power supply. The control of these devices is made by a computer through a GPIB connection and a specific software that has been developed. The change of heating power of the control-heater before, during and after measurements is a direct measure for the excess enthalpy H^E . The calibration of the measurement system is made by simulating an exothermic mixing process by a calibration resistor.

Knowing the volumetric flow rates delivered, the molar masses and the densities of the pure compounds, the mole fractions of the mixtures obtained in the mixing coil can be calculated. The maximum absolute uncertainty of mole fraction at equimolar composition is ± 0.0008 . Densities of pure liquids are determined interpolating density data obtained from Riddick et al.⁶ at the measured temperature of delivery. Table 1 shows agreement between our estimated density data and literature data. The temperature of pure liquids is measured by a calibrated standard PRT-100 inserted in the pump

body, using as indicator an a/c resistance bridge (ASL model F250) resolving 1 mK in the reading of temperature and estimating an overall uncertainty of ± 10 mK. Mixtures of different compositions are studied by changing the ratio of flow rates and in this way the dependence of H^E on mole fraction can be determined. The estimated relative uncertainty of the determined $H^E/\text{J}\cdot\text{mol}^{-1}$ is $\pm 0.01\cdot H^E$.

To check the above apparatus we have used two systems traditionally considered as test systems for calorimetric techniques: cyclohexane + hexane and benzene + cyclohexane at 298.15 K. The first system, cyclohexane (1) + hexane (2), was measured by Marsh and Stokes¹² with an isothermal displacement microcalorimeter. This work presents the most complete amount of data and is taken as reference system. However, we have considered also the comparison with data from Murakami et al.¹³, which used a continuous dilution calorimeter, and from Gmehling et al.¹⁴, whose data were taken with an isothermal flow calorimeter of the same characteristics as ours. Table 2 presents the results found in the references for this system and the results obtained in this work, as well as the maximum deviation of the excess enthalpy respect the calculated value, $\max|\Delta H^E|$, and the standard deviation, $\text{rms } \Delta H^E$. The Redlich-Kister equation is used for the adjustment of each data set, with the number of parameters employed by each author:

$$H^E = x \cdot (1-x) \cdot \sum_{i=1}^n A_i \cdot (2x-1)^{i-1} \quad (1)$$

The comparison of the calculated values of the Redlich-Kister coefficients and scattering parameters in this work with those from the literature, shows that our values are similar with regard to the same type of isothermal flow calorimeter and more scattered in relation with isothermal displacement microcalorimeter and continuous dilution calorimeter.

A comparison between experimental data of this work and those from the literature is presented in Figure 3. For each data set, the calculated enthalpy H^E_{calc} has been determined using the Redlich-Kister coefficients of the correlation determined by Marsh and Stokes¹². The scattering of experimental data obtained in this work is within the $\pm 1\%$ of the calculated enthalpy corresponding to each mole fraction in the central range of composition.

The second test system is the binary mixture benzene (1) + cyclohexane (2) at 298.15 K. In a similar way as for the former system, comparison between the data of Ewing et al.¹⁵, Lundberg¹⁶, Savini et al.¹⁷ and this work is reported in Table 2 and Figure 3, being the coefficients of Ewing et al.¹⁵ used to obtain the calculated enthalpy H^E_{calc} for each data set. In this case, also the agreement between this work and literature data is within the $\pm 1\%$ of the calculated enthalpy.

Results and Discussion

The experimental excess molar enthalpies obtained in this work for the binary mixtures DBE (1) + cyclohexane (2), DBE (1) + 1-butanol (3) and cyclohexane (2) + 1-butanol (3) at 298.15 K are listed in Table 3.

For binary systems, there are several models and empirical equations proposed to fit the H^E measurements. One of them, the Redlich-Kister equation, is given by eq. 1, in which the A_i coefficients are determined by the unweighted least-squares method. Another equation very employed is the modified Margules eq. 2, proposed by Abbot and van Ness¹⁸

$$H^E = x_1 x_2 RT \left[a_{21} x_1 + a_{12} x_2 - \frac{x_1 x_2 b_{12} b_{21}}{b_{21} x_1 + b_{12} x_2 + x_1 x_2 (c_{21} x_1 + c_{12} x_2)} \right] \quad (2)$$

Binary systems have also been correlated using NRTL model, which expression for the molar excess enthalpy is indicated by the eq. 3:

$$H^E = -RT \sum_{i=1}^n x_i \eta_i \quad (3)$$

where x_i is the mole fraction of the component i and η_i is given by

$$\eta_i = \frac{\sum_{k=1}^p x_k \tau_{ki} G_{ki} \left[\alpha \left(\tau_{ki} - \left(\frac{\sum_{n=1}^p x_n \tau_{ni} G_{ni}}{\sum_{l=1}^n x_l G_{li}} \right) \right) - 1 \right]}{\sum_{l=1}^p x_l G_{li}}$$

$$G_{ji} = \exp(-\alpha \tau_{ij})$$

The NRTL theory assumes that the excess molar enthalpy of a multicomponent mixture depends only on the binary parameters. In this work, the non-randomness parameter α was assigned 0.3 for the NRTL model of two parameters, and as an adjustable parameter for the NRTL model of three parameters.

For the UNIQUAC model, the molar excess enthalpy is given by

$$H^E = \sum_{i=1}^n q_i x_i \frac{\sum_{j=1}^n g_j \Delta u_{ji} \tau_{ji}}{\sum_{j=1}^n g_j \tau_{ij}} \quad (4)$$

where $g_i = \frac{q_i x_i}{\sum_j q_j x_j}$ and q_i the molecular surface area, obtained as the sum of the contributions of the functional groups presents in the compound.

Results of data correlation for the binary systems are summarized in Table 4. A plot of the experimental and correlated data is shown in Figure 4.

The ternary mixtures DBE (1) + cyclohexane (2) + 1-butanol (3) at 298.15 K were formed by adding the cyclohexane (2) to binary mixtures of fixed composition of DBE (1) + 1-butanol (3). Four different starting binaries were used, with values of the ratio x_1/x_3 of 0.2498, 0.6669, 1.4988 and 4.0531 respectively. The experimental excess molar enthalpies listed in Table 5 are determined by eq. 5, using the smooth representation of H^E_{13} by Redlich-Kister:

$$H^E_{123} = H^E_{2+13} + (1 - x_2)H^E_{13} \quad (5)$$

Experimental results for the excess molar enthalpy H^E_{123} are shown in Figure 5.

The following equation was used to fit the H^E measurements

$$H^E_{123} = H^E_{12} + H^E_{13} + H^E_{23} + x_1 x_2 x_3 \Delta H^E_{123} \quad (6)$$

with

$$\Delta H^E_{123} = B_0 + B_1 x_1 + B_2 x_2 + B_3 x_1^2 + B_4 x_2^2 + B_5 x_1 x_2 + B_6 x_1^3 + B_7 x_2^3 \quad (7)$$

where the parameters B_i were determined by the unweighted least-squares method. Another alternative for the calculation of ΔH^E_{123} is eq. 8

$$\Delta H_{123}^E = B_1x_1 + B_2x_2 + B_3x_3 \quad (8)$$

Table 6 presents the summary of correlation results obtained for the ternary system with eq. 6, 7, 8, and NRTL and UNIQUAC models. Table 6 also shows the prediction results obtained for the ternary system using the parameters of the binary systems.

The excess molar enthalpy of the system DBE (1) + cyclohexane (2) at 298.15 K shows endothermic behaviour ($H^E > 0$) in the whole range of composition. The best fit of experimental data is obtained with the Redlich-Kister equation with a root mean square deviation, rms ΔH^E , of 0.5 J·mol⁻¹ and a maximum value of the absolute deviation, max $|\Delta H^E|$, of 1.2 J·mol⁻¹. Deviations obtained with NRTL and UNIQUAC models are higher. The maximum value of the molar excess enthalpy is 267 J·mol⁻¹, obtained at a mole fraction of DBE about 0.40. Comparison with data for the same binary system and temperature reported by Marongiu et al.¹⁹ shows that our H^E data are considerably higher, the difference being 15 % in the central range of composition, even when the calorimeter was re-calibrated and the binary system measured three times.

The binary system DBE (1) + 1-butanol (3) measured at 298.15 K is also endothermic. The maximum value of H^E is 892 J·mol⁻¹, at a mole fraction of DBE about 0.60. Modified Margules equation gives the best fit with a root mean square deviation, rms ΔH^E , of 2.4 J·mol⁻¹ at 298.15 K. The maximum values of the absolute deviation, max $|\Delta H^E|$, is 4.0 J·mol⁻¹. The remaining models give worse fits. Our H^E data agree to within 1,5 % in the central range of composition with the measurements for the same system and temperature obtained from Kammerer and Lichtenthaler.²⁰

The third binary system which has been studied is the system cyclohexane (2) + 1-butanol (3) at 298.15 K, showing also an endothermic behaviour. The maximum value of the molar excess enthalpy of the mixture is 635 J·mol⁻¹. The mole fraction of cyclohexane at this maximum is approximately 0.65. The experimental data have been fitted by Redlich-Kister, modified Margules, NRTL and UNIQUAC models, being the Modified Margules equation the best fit with a root mean square deviation, rms ΔH^E , of 2.6 J·mol⁻¹. The maximum value of the absolute deviation, max $|\Delta H^E|$, is 6.0 J·mol⁻¹. Comparison with

1 data for the same binary system and temperature reported by Löwen and Schulz²¹ shows that our H^E data
2 agree to within 1.7 % in the central range of composition.
3

4
5 Concerning the measured ternary system, the best fit of experimental data is obtained with eq. 6 and 7.
6
7 The root mean square deviation, rms ΔH^E , is of 5.0 J·mol⁻¹, and the maximum value of the absolute
8 deviation, $\max|\Delta H^E|$, is 13.4 J·mol⁻¹ which means about 5 % over the experimental molar excess
9 enthalpy at this point. The results obtained with the other models are rather worse. This system shows an
10 endothermic behaviour in the whole range of composition. The maximum value of H^E is 890 J·mol⁻¹. No
11 data of the same system were found in the literature for comparison. Both NRTL and UNIQUAC models
12 present high root mean square deviation values in the prediction of data.
13
14
15
16
17
18
19
20

21 **Conclusions**

22
23
24 Isothermal excess enthalpies at 298.15 K for the ternary system dibutyl ether (DBE) + cyclohexane +
25 1-butanol and its constituent binary systems were determined by using an isothermal flow calorimeter.
26
27 All the binary systems show positive values and strongly asymmetric H^E behavior in the measured
28 temperature. The measured excess enthalpies data were correlated well with various common H^E
29 models.
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

FIGURE CAPTIONS




Figure 1. Quasi-isothermal flow calorimeter. Schematic diagram.






Figure 2. Mixing flow cell. Schematic diagram.



(a)




(b)


Figure 3. Comparison of residuals between the measured excess enthalpy, H^E , and the calculated excess enthalpy, H^E_{calc} , at the temperature of 298.15 K, for the following systems: (a) Cyclohexane (1) + Hexane (2), O, This work; x, Marsh and Stokes¹²; Δ , Murakami and Benson¹³; \square , Gmehling and

1 Meents¹⁴. (b) Benzene (1) + Cyclohexane (2), O, This work; x, Ewing et al.¹⁵; □ Lundberg¹⁶; Δ, Savini
2
3 et al.¹⁷.
4
5


6 In Figure 3(a), the calculated enthalpy H^E_{calc} has been determined using the Redlich-Kister coefficients
7
8 of the correlation determined by Marsh and Stokes⁷ for each data set, while those from Ewing et al.¹⁵
9
10 have been used in Figure 3(b). The dotted line shows the $\pm 1\%$ deviation from the calculated enthalpy
11
12 corresponding to each mole fraction.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



(a)




(b)



(c)

Figure 4. Excess molar enthalpy H^E at 298.15 K for (a) DBE (1) + cyclohexane (2); (b) DBE (1) + 1-butanol (3); (c) cyclohexane (2) + 1-butanol (3). Experimental results: (\square); (—) calculated with eq. 1 (Figure a) and eq. 2 (Figures b and c) using parameters of Table 4.



35 **Figure 5.** Contours for constant values of H^E_{123} for DBE(1) + cyclohexane (2) + 1-butanol (3) at 298.15

36
37 K.
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

TABLES.

Table 1. Comparison of density data of pure compounds, cyclohexane, hexane, benzene, DBE and 1-butanol, at 298.15 K.

	cyclohexane		hexane		benzene		DBE		1-butanol	
	this work	lit.	this work	lit.						
$\rho/\text{g}\cdot\text{cm}^{-3}$	0.77389 ^a	0.7738 ^b	0.65484 ^a	0.6551 ^c	0.87360 ^a	0.8736 ^d	0.76410 ^a	0.76411 ^e	0.80575 ^a	0.80561 ^f

^aRiddick et al.⁶. ^bRodríguez et al.⁷. ^cIglesias et al.⁸. ^dIglesias et al.⁹. ^eJiménez et al.¹⁰. ^fRezanova et al.¹¹.

Table 2. Comparison of H^E data for the binary systems cyclohexane + hexane and benzene + cyclohexane, at 298.15 K. Parameters A_i of Redlich-Kister equation, standard deviation rms ΔH^E and maximum excess enthalpy deviation max. $|\Delta H^E|$.

Calorimeter technique	A_1	A_2	A_3	A_4	rms ΔH^E /J·mol ⁻¹	max. $ \Delta H^E $ /J·mol ⁻¹
Cyclohexane (1) + Hexane (2)						
Isothermal displacement calorimeter. ^a	864.7	249.5	99.1	33.2	0.1	0.2
Continuous dilution calorimeter. ^b	863.5	248.8	104.5	31.1	0.2	0.4
Isothermal flow calorimeter. ^c	861.6	264.4	103.8		0.8	1.6
Isothermal flow calorimeter. ^d	867.2	245.3	63.8	23.7	0.8	1.4
Benzene (1) + Cyclohexane (2)						
Isothermal displacement calorimeter. ^e	3197.1	175.4	121.5	28.5	0.2	0.5
Continuous dilution calorimeter. ^f	3196.7	153.9	123.0		2.0	2.7
Continuous dilution calorimeter. ^g	3190.4	163.5	112.5	36.7	0.3	0.6
Isothermal flow calorimeter. ^d	3201.3	171.8	151.5		0.9	2.3

^aMarsh and Stokes¹². ^bMurakami and Benson¹³. ^cGmehling and Meents¹⁴. ^eEwing et al.¹⁵. ^fLundberg¹⁶.
^gSavini et al.¹⁷. ^dThis work

Table 3. Experimental excess molar enthalpies of binary systems DBE (1)+ cyclohexane (2), DBE (1) + 1-butanol (3) and cyclohexane (2) + 1-butanol (3) at 298.15 K.

x	$H^E/\text{J}\cdot\text{mol}^{-1}$	x	$H^E/\text{J}\cdot\text{mol}^{-1}$	x	$H^E/\text{J}\cdot\text{mol}^{-1}$	x	$H^E/\text{J}\cdot\text{mol}^{-1}$
x DBE + $(1-x)$ cyclohexane							
0.0499	75.2	0.2999	255.4	0.5495	248.1	0.7990	136.7
0.0997	134.1	0.3500	264.9	0.6000	232.6	0.8500	104.1
0.1502	180.4	0.3996	268.6	0.6491	213.9	0.9001	69.3
0.2002	214.5	0.4497	266.4	0.6992	191.1	0.9507	32.8
0.2497	238.5	0.4990	259.6	0.7501	164.1		
x DBE + $(1-x)$ 1-butanol							
0.0501	119.4	0.3004	624.6	0.5506	879.8	0.8000	772.8
0.1002	236.0	0.3498	698.4	0.5998	892.4	0.8501	679.1
0.1497	344.5	0.4006	762.1	0.6505	890.6	0.8994	543.1
0.2003	447.6	0.4507	814.0	0.7000	872.6	0.9493	330.1
0.2503	540.2	0.4997	853.2	0.7509	833.2		
x cyclohexane + $(1-x)$ 1-butanol							
0.0503	68.3	0.2997	407.6	0.5501	613.4	0.7996	567.1
0.0998	144.8	0.3500	461.6	0.5997	629.5	0.8500	513.0
0.1502	217.4	0.4002	510.5	0.6501	634.6	0.9000	438.5
0.1998	286.8	0.4503	552.5	0.7002	626.7	0.9496	334.4
0.2502	349.6	0.5003	587.3	0.7501	604.7		

Table 4. Summary of parameters for the representation of H^E by Redlich-Kister, Modified Margules equation, NRTL (2 and 3 parameters) and UNIQUAC models, for binary systems DBE (1) + cyclohexane (2), DBE (1) + 1-butanol (3) and cyclohexane (2) + 1-butanol (3) at 298.15 K.

Binary Systems ^a	Correlation				
	Redlich-Kister	Modified Margules	NRTL	NRTL(3p)	UNIQUAC
DBE (1)+ cyclohexane (2)					
A_0	1037.1	0.979	-0.2065	0.1153	-210.2
A_1	-362.2	0.277	0.8622	0.5691	467.4
A_2	166.2	0.328			
A_3	-139.2	117.8			
A_4	-29.0	2.6			
A_5		-106.4			
α_{12}			0.30	0.82	
rms $\Delta H^E/\text{J}\cdot\text{mol}^{-1}$	0.5	0.6	2.9	1.5	1.4
Max $ \Delta H^E /\text{J}\cdot\text{mol}^{-1}$	1.2	1.5	5.0	3.8	3.5
Max $(\Delta H^E /H^E)$	3.6 %	4.5 %	5.7 %	11.5 %	10.5 %
DBE (1) + 1-butanol (3)					
A_0	3414.8	1.011	2.3362	2.4081	2160.1
A_1	1259.1	1.037	0.6854	0.8333	-629.7
A_2	741.4	-256347			
A_3	1310.7	-0.5147			
A_4	1021.8	72852			
A_5		201903			
α_{13}			0.30	0.31	
rms $\Delta H^E/\text{J}\cdot\text{mol}^{-1}$	2.9	2.4	14.3	11.9	43.2
Max $ \Delta H^E /\text{J}\cdot\text{mol}^{-1}$	4.9	4.0	29.0	21.8	84.5
Max $(\Delta H^E /H^E)$	3.5 %	2.6 %	8.8 %	9.7 %	23.1 %
cyclohexane (2)+1-butanol(3)					
A_0	2339.6	0.6024	1.9691	2.1696	1960.8
A_1	1371.5	8.2647	0.0910	0.6426	-588.4
A_2	1046.8	410.0			
A_3	-1202.8	7.181			
A_4	-2123.2	16.9			
A_5	3793.8	4.5			
A_6	4414.9				

α_{23}			0.30	0.42	
rms $\Delta H^E/\text{J}\cdot\text{mol}^{-1}$	6.5	2.6	36.3	24.8	37.2
Max $ \Delta H^E /\text{J}\cdot\text{mol}^{-1}$	13.6	6.0	111.2	75.7	116.0
Max ($ \Delta H^E /H^E$)	8.0 %	8.9 %	33.2 %	26.2 %	34.7 %

^a Equivalence between parameters: NRTL $A_0=\tau_{12}$ and $A_1=\tau_{21}$; UNIQUAC $A_0=\Delta u_{12}$; $A_1=\Delta u_{21}$;
Modified Margules $A_0=a_{12}$; $A_1=a_{21}$; $A_2=b_{12}$; $A_3=b_{21}$; $A_4=c_{12}$; $A_5=c_{21}$

For Review - Confidential - ACS

Table 5. Experimental excess molar enthalpies H^E_{2+13} at 298.15 K for the addition of cyclohexane to (DBE (1) + 1-butanol (3)) to form x_1 DBE + x_2 cyclohexane + $(1-x_1-x_2)$ 1-butanol, and values of H^E_{123} calculated from eq. 5, using the smooth representation of H^E_{13} by Redlich-Kister equation with parameters given in Table 4.

x_2	$H^E_{2+13}/\text{J}\cdot\text{mol}^{-1}$	$H^E_{123}/\text{J}\cdot\text{mol}^{-1}$	x_2	$H^E_{2+13}/\text{J}\cdot\text{mol}^{-1}$	$H^E_{123}/\text{J}\cdot\text{mol}^{-1}$
		$x_1/x_3=0.2498$	$H^E_{13}/\text{J}\cdot\text{mol}^{-1}=443.8$		
0.9000	394.4	395.0	0.4004	475.1	579.0
0.8002	504.4	508.6	0.2998	387.6	549.7
0.6996	557.4	571.3	0.2003	277.7	514.2
0.6003	565.3	597.4	0.1001	144.2	474.2
0.5002	536.1	597.6			
		$x_1/x_3=0.6669$	$H^E_{13}/\text{J}\cdot\text{mol}^{-1}=764.2$		
0.8999	350.8	351.6	0.4006	422.9	592.1
0.7997	446.5	453.0	0.3007	348.0	614.8
0.6998	491.2	512.9	0.2003	250.1	646.2
0.6002	498.3	549.3	0.0997	130.0	691.4
0.5002	473.6	572.4			
		$x_1/x_3=1.4988$	$H^E_{13}/\text{J}\cdot\text{mol}^{-1}=890.0$		
0.9002	309.0	309.8	0.4001	369.5	554.9
0.8000	395.9	402.5	0.2998	303.5	600.9
0.6997	433.8	456.5	0.2005	216.8	663.4
0.5998	438.3	492.4	0.1001	112.8	755.5
0.4998	415.3	521.9			
		$x_1/x_3=4.0531$	$H^E_{13}/\text{J}\cdot\text{mol}^{-1}=773.3$		
0.8998	256.9	257.5	0.4005	322.0	460.4
0.8006	342.6	347.1	0.2999	263.1	492.5
0.7002	380.0	395.8	0.1997	186.6	544.6
0.6003	383.8	422.1	0.0995	97.1	632.5
0.5009	363.7	440.8			

Table 6. Summary of the data reduction and prediction results obtained for the ternary system DBE (1)

+ cyclohexane (2) + 1-butanol (3) at 298.15 K.

CORRELATION ^a	ΔH^E_{123} (eq. 7)	ΔH^E_{123} (eq. 8)	NRTL	UNIQUAC
B_0	2586.7		-0.3508	1294.6
B_1	-12380.7	-1936.1	0.4952	-711.7
B_2	-15976.6	-4187.3	1.7410	1416.8
B_3	17886.7	911.9	0.6327	-389.5
B_4	39625.0		3.7942	2869.9
B_5	7477.6		0.1853	-825.9
B_6	-12474.9			
B_7	-41723.8			
α_{12}			0.30	
α_{13}			0.30	
α_{23}			0.30	
rms $\Delta H^E/\text{J}\cdot\text{mol}^{-1}$	5.0	11.9	49.9	55.2
Max $ \Delta H^E /\text{J}\cdot\text{mol}^{-1}$	13.4	27.5	94.1	178.5
Max $(\Delta H^E /H^E)$	5.2 %	8.9 %	20.0 %	44.9 %
PREDICTION ^a			NRTL	UNIQUAC
B_0			-0.2065	-210.2
B_1			0.8622	467.4
B_2			2.3362	2160.1
B_3			0.6854	-629.7
B_4			1.9691	1960.8
B_5			0.0910	-588.4
α_{12}			0.30	
α_{13}			0.30	
α_{23}			0.30	
rms $\Delta H^E/\text{J}\cdot\text{mol}^{-1}$			207.0	169.8
Max $ \Delta H^E /\text{J}\cdot\text{mol}^{-1}$			356.2	306.9
Max $(\Delta H^E /H^E)$			64.2 %	55.3 %

^a Equivalence between parameters: NRTL $B_0=\tau_{12}$; $B_1=\tau_{21}$; $B_2=\tau_{13}$; $B_3=\tau_{31}$; $B_4=\tau_{23}$; $B_5=\tau_{32}$; UNIQUAC $B_0=\Delta u_{12}$; $B_1=\Delta u_{21}$; $B_2=\Delta u_{13}$; $B_3=\Delta u_{31}$; $B_4=\Delta u_{23}$; $B_5=\Delta u_{32}$;

Literature Cited

- 1
2
3
4 (1) Redlich, O.; Kister, A. T. Algebraic Representation of Thermodynamic Properties and the
5
6 Classification of Solutions. *Ind. & Eng. Chem.* **1948**, *40*, 345-348.
- 7
8
9 (2) Renon, H.; Prausnitz, J. M. Local Compositions in Thermodynamic Excess Functions for Liquid
10
11 Mixtures. *AIChE Journal* **1968**, *14*, 135-144.
- 12
13
14 (3) Abrams, D. S.; Prausnitz, J. M. Statistical Thermodynamics of Liquid Mixtures: A New
15
16 Expression for the Excess Gibbs Energy of Partly or Completely Miscible Systems. *AIChE*
17
18 *Journal* **1975**, *21*, 116-128.
- 19
20
21 (4) Alonso, C.; Montero, E. A.; Chamorro, C. R.; Segovia, J. J.; Martín, M. C.; Villamañán, M. A.
22
23 Excess Enthalpies of Binary and Ternary Mixtures Containing tert-Amyl Methyl Ether (TAME),
24
25 tert-Amyl Alcohol (TAOH) and Hexane at 298.15 and 313.15 K. *Fluid Phase Equilib.* **2004**,
26
27 *217*, 145-155.
- 28
29
30 (5) Alonso, C. Ph. D. Thesis, University of Valladolid, Spain, 2001.
- 31
32
33 (6) Riddick, J. A.; Bunger, W. B.; Sakano, T. K. *Organic Solvents, Physical Properties and Methods*
34
35 *of Purification*; Wiley; New York, 1986.
- 36
37
38 (7) Rodríguez, A., Canosa, J., Tojo, J. Physical properties of the binary mixtures (diethyl carbonate
39
40 + hexane, heptane, octane and cyclohexane) from T=293.15 K to T=313.15K. *J. Chem.*
41
42 *Thermodyn.* **2003**, *35*, 1321-1333.
- 43
44
45 (8) Iglesias, M., Orge, B., Piñeiro, M. M., Marino, G., Tojo, J. Volumetric properties prediction by
46
47 cubil EOS for non-ideal mixtures: application to the ternary system acetone + methanol + n-
48
49 hexane. *Thermochim. Acta* **1999**, *328*, 265-275.
- 50
51
52 (9) Iglesias, M., Piñeiro, M. M., Marino, G., Orge, B., Domínguez, M., Tojo, J. Thermodynamic
53
54 properties of the mixture benzene + cyclohexane + 2-methyl-2-butanol at the temperature 298.15
55
56
57
58
59
60

1 K: excess molar volumes prediction by application of cubic equations state. *Fluid Phase Equilib.*

2
3 **1999**, *154*, 123-138.

4
5
6 (10) Jiménez, E., Segade, L., Franjo, C., Casas, H., Legido, J.L., Paz Andrade, M.I. Viscosity
7 deviations of ternary mixtures di-n-butyl ether + 1-propanol + n-octane at several temperatures.

8
9
10 *Fluid Phase Equilib.* **1998**, *149*, 339-358.

11
12
13 (11) Rezanova, E. N., Kammerer, K., Lichtenthaler, R.N. Excess enthalpies and volumes of ternary
14 mixtures containing 1-propanol or 1-butanol, an ether (diisopropyl ether or dibutyl ether) and
15 heptane. *J. Chem. Eng. Data* **2000**, *45*, 124-130.

16
17
18 (12) Marsh, K. N.; Stokes, R. H. Enthalpies of Mixing of n-Hexane + Cyclohexane at 25°C. *J. Chem.*
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Thermodyn. **1969**, *1*, 223-225.

(13) Murakami, S.; Benson, G. C. *Int. Data Series (A)* 1974 (1), 19.

(14) Gmehling, J.; Meents, B. *Int. Data Series (A)* 1992 (3), 144.

(15) Ewing, M. B.; Marsh, K. N.; Stokes, R. H.; Tuxford, C. W. *Int. Data Series (A)* 1973, 23.

(16) Lundberg, G. W. *Int. Data Series (A)* 1974 (3), 225.

(17) Savini, G. C.; Winterhalter, D. R.; Kovach, L. H.; Van Ness, H. C. *Int. Data Series (A)* 1974 (3),
160.

(18) Abbott, M. M; Van Ness, H. C. Vapor –Liquid Equilibrium: Part III. Data Reduction with
Precise Expressions for G^E . *AIChE Journal* **1975**, *21*, 62-71.

(19) Marongiu, B; Dermini, S.; Lepori, L.; Matteoli, E.; Kehiaian, H.V. Thermodynamics of Binary
Mixtures Containing Ethers or Acetals. 1. Excess Enthalpies of Linear Ethers or Acetals +
Heptane or + Cyclohexane Mixtures. *J. Chem. Eng. Data* **1988**, *33*, 118-122.

- 1 (20) Kammerer, K.; Lichtenthaler, R. N. Excess Properties of Binary Alkanol-Ether Mixtures and the
2 Application of the ERAS Model. *Thermochim. Acta* **1998**, *310*, 61-67.
3
4
5
6 (21) Löwen, B.; Schulz, S. Excess Molar Enthalpies of Cyclohexane + n-Alcohols at 283.15, 298.15,
7
8 323.15 and 363.15 K and at a Pressure of 0.4 MPa. *Thermochim. Acta* **1995**, *265*, 63-71.
9

10
11 Support for this work came from the Dirección General de Investigación (DGI), Ministerio de
12 Educación y Ciencia, Spain, Projects ENE2006-12620 and ENE2006-13349, and from the Consejería de
13 Educación, Junta de Castilla y León, Spain, Project BU015A06.
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60