



# Stefano Marcaccini: a pioneer in isocyanide chemistry



Ana G. Neo<sup>1</sup> · José Luis Ramiro<sup>1</sup> · María García-Valverde<sup>2</sup> · Jesús Díaz<sup>1</sup> · Carlos F. Marcos<sup>1</sup>

Received: 23 January 2023 / Accepted: 20 March 2023  
© The Author(s) 2023

## Abstract

Stefano Marcaccini was one of the pioneers in the use of isocyanide-based multicomponent reactions in organic synthesis. Throughout his career at the University of Florence he explored many different faces of isocyanide chemistry, especially those geared towards the synthesis of biologically relevant heterocycles. His work inspired many researchers who contributed to other important developments in the field of multicomponent reactions and created a school of synthetic chemists that continues today. In this manuscript we intend to review the articles on isocyanide multicomponent reactions published by Dr. Marcaccini and analyse their influence on the following works by other researchers. With this, we hope to highlight the immense contribution of Stefano Marcaccini to the development of isocyanide chemistry and modern organic synthesis as well as the influence of his research on future generations. We believe that this review will not only be a well-deserved tribute to the figure of Stefano Marcaccini, but will also serve as a useful inspiration for chemists working in this field.

## Graphical abstract



**Keywords** Isocyanides · Multicomponent reactions · Heterocycles · Organic synthesis · Cycloadditions

## Introduction

Stefano Marcaccini was born in Prato (Florence) on the 31 of January of 1956. He studied at the University of Florence, where he got a PhD in Organic Chemistry in 1982 under the supervision of Professor Valerio Parrini. Between 1994 and 1995, he held a position as associate professor of Heterocyclic Chemistry at the University of Siena. He then returned to the University of Florence, where he spent most of his career, conducting outstanding research on the synthesis of

✉ Carlos F. Marcos  
cfernan@unex.es

<sup>1</sup> Laboratory of Bioorganic Chemistry & Membrane Biophysics (L.O.B.O.), Universidad de Extremadura, 10003 Cáceres, Spain

<sup>2</sup> Departamento de Química, Facultad de Ciencias, Universidad de Burgos, 09001 Burgos, Spain

heterocycles that led to novel methods based primarily on isocyanide reactions.

He became a world recognised expert on the chemistry of isocyanides, having published a hundred of research papers on this field. His contribution has an unquestionable impact on the current state of the art of multicomponent reactions. He mentored many students, and he was appreciated, not only for his quality as educator and his extraordinary chemical intuition, but especially for his exceptional generosity and kindness. His work inspired many researchers and sparked the creation of a school of synthetic chemists that is still very active today.

Stefano Marcaccini died at Prato on the 1st of October 2012. He is remembered as a remarkably gifted chemist and an excellent person.



## Synthesis of heterocycles from bifunctional isocyanides

### Isocyanacetate derivatives

In the mid-80s, Marcaccini's work was focused on heterocyclic chemistry. Thus, with the aim of developing novel ways to achieve heterocyclic scaffolds he turned the spotlight into isocyanide chemistry, specifically isocyanacetate

derivatives (**1**). He reasoned that the acidic position  $\alpha$  to the isocyanide could be used to cyclise onto suitable functional groups introduced with other reagents. He was probably inspired by the seminal research of Schöllkopf on the coupling of isocyanoacetate (**1**) with acyl chlorides (**6**) [1, 2], and of Van Leusen, on *p*-toluenesulfonylmethyl isocyanide (TOSMIC) chemistry [3].



Thus, he reacted 2-isocyanoacetate (**1**) with sulphur electrophiles, such as sulphenyl chlorides (**2**), generating a reactive intermediate (**3**), which would undergo an intramolecular cyclisation in the presence of a base. He and his co-workers synthesised different heterocyclic systems using this strategy. For example, in his first work with isocyanides, 2-arylthio-5-alkoxyoxazoles (**5**) were constructed in almost quantitative yields in a one pot process (Scheme 1) [4]. A few years later, Marcaccini and Torroba expanded this study by acylating oxazole (**5**) to afford trisubstituted oxazoles (**7**; Scheme 1) [5]. Most likely, these reactions proceed through nitrilium ylide intermediates (**4**), which have ever since been revealed as a common and very useful feature of the chemistry of isocyanoacetate derivatives offering a broad variety of a common and very useful feature applications in the multicomponent synthesis of heterocycles [6].


Similarly, dioxazolylsulfides (**11**) were prepared from alkyl isocyanoacetates (**1**) and sulphur dichloride (**8**). A 2:1 molar ratio was used in this case to favour the formation of a labile intermediate (**10**), which easily cyclised in presence of a base leading to bisoxazolyl sulfane (**11**) in high yields (Scheme 2) [7].

Additionally, the use of alkyl isocyanoacetates (**12**) and  $S_2Cl_2$  (**13**) unexpectedly resulted in a convenient synthesis of fused heterocyclic cores, such as thiazolo[5,4-*d*]-thiazoles (**21**; Scheme 3) [8]. This molecular system has been recognised as an important motif for molecular optic-electronic [9] and photovoltaic applications [10]. The mechanism of the reaction appears to involve a dimerisation of intermediate (**16**), which was confirmed as the same thiazolo [5,4-*d*] thiazole (**21**) that can be obtained by reacting  $S_2Cl_2$  (**8**) with isocyanoacetate (**12**) in a 1:1 molar ratio.


Other complex heterocyclic systems, such as 6-arylthio-8-ethoxycarbonyl-4-ethoxycarbonylmethylaminoimidazo[5,1-*b*][1,3,5]thiadiazine-2-thiones (**28**) could be obtained from alkyl isocyanoacetates (**12**) and arylsulphenyl thiocyanates (**22**). The intermediate salt (**27**) was treated with acid to afford thione (**28**), which could be further methylated to give the corresponding  $SCH_3$  derivatives (**30**; Scheme 4) [11].

Inspired by the general idea of Marcaccini's seminal works, different strategies have recently been developed to elicit the coupling of isocyanides with sulphur electrophiles. For example, Mampuy et al. synthesised secondary thiocarbamates (**33**) from isocyanides (**31**) and thiosulfonates (**32**) in a transition metal-free protocol enabled by the catalytic

**Scheme 1** Synthesis of substituted oxazoles**Scheme 2** Synthesis of 5,5'-dialkoxy-2,2'-dioxazolyldisulfides**Scheme 3** Synthesis of thiazolo[5,4-*d*]-thiazoles



**Scheme 4** Synthesis of 6-arylthio-8-ethoxycarbonyl-4-ethoxycarbonylmethylaminoimidazo[5,1-*b*][1,3,5]thiadiazine-2-thiones




**Scheme 5** Secondary thiocarbamate synthesis from isocyanides

electron transfer role played by inexpensive NaI (Scheme 5) [12].

Using a novel perspective, Guan's group employed an organic electrochemistry approach for the activation of isocyanides (31) with sulphur electrophiles. In this approach, the reaction of isocyanides (31) and thiols (35) afforded a wide range of imino sulphide ethers (37), which are important as pharmaceuticals and as key intermediates in sugar chemistry (Scheme 6) [13].

An interesting related approach has been reported by Sun and co-workers, who reacted disulphides (38) with isocyanides (31) in the presence of TEMPO and *N*-halosuccinimides (39) to give insertion products that can further incorporate a wide range of nucleophiles (41) [14]. This sequence led to the multicomponent construction of different molecular scaffolds (42) in mild conditions (Scheme 7).

Isocyanide insertion into S–S bonds can be also used in the construction of heterocycles. Thus, Marcos et al. developed a straightforward method for the synthesis of 2-amino-benzothiazoles (48) through an iodine-catalysed insertion of





**Scheme 6** Electrochemical synthesis of imino-sulphide ethers

isocyanides into the S–S bond of benzodithiazole 2-oxides (**45**), with concomitant extrusion of sulphur monoxide (Scheme 8) [15].

The insertion of isocyanides into N–Cl bonds is also possible. Marcaccini's group exploited the chemistry of isocyanides to synthesise different unusual molecules not


easily attainable by other procedures, highlighting the great potential of this functional group [16]. For example, they easily obtained sulfonylguanidines (**54**) through the reaction of isocyanides (**31**) with anilines (**49**) and chloramine T (**50**) under phase transfer-catalysed conditions at room temperature (Scheme 9) [17].

**Scheme 7** Activated disulphides in reactions with isocyanides and nucleophiles



**Scheme 8** Marcos' synthesis of 2-amino-benzothiazoles

**Scheme 9** Synthesis of sulfonylguanidines



The authors argue that the formation of *N*-chloroamines (**51**) is the crucial step in this synthesis. Specifically, they propose that the reaction takes place by the *N*-chlorination of aromatic amines (**49**) to give *N*-chloroanilines (**51**), which react with isocyanides (**31**) to give the  $\alpha$ -adducts (**53**). These then react with sodium tosylamide (**52**) to form the sulfonylguanidines (**54**).

The use of bifunctional amines may facilitate post-condensation transformations leading to heterocyclic products. Therefore, when the reaction was carried out with methyl anthranilate (**55**) the resulting intermediate sulfonylguanidine (**57**) underwent a heating mediated cyclisation to readily give quinazoline derivatives (**58**; Scheme 10) [18].

### Isocyanacetamide derivatives

Isocyanacetamides (**59**), the corresponding amide derivatives of alkyl isocyanacetates (**1**), are also powerful reagents for heterocyclic synthesis with exceptional characteristics [19].

Marcaccini et al., in parallel with their research on 2-isocyanacetates (**1**), pioneered the reaction between isocyanacetamides (**59**) and sulphur chloride derivatives, such as aryl sulfonyl chlorides (**60**), to afford mesoionic heterocycles (**62**; Scheme 11) [20]. As in similar reactions with isocyanacetates (**1**), the product is obtained by a base mediated cyclisation of the  $\alpha$ -addition intermediate (**61**).

Following this, García-Valverde and Marcaccini developed a novel regioselective and experimentally simple synthesis of iminohydantoins from isocyanacetamides (**63**) and chloroamines (**56**). 2-Iminohydantoins (**66**) and 4-iminohydantoins (**70**) were selectively obtained starting from *N*-aryl (**63**) or *N*-alkylisocyanacetamides (**67**), respectively (Scheme 12) [21]. According to the mechanism proposed, regioselectivity is controlled by the electron density of the amide nitrogen. Aryl substituents generate lower electron


density at the amide nitrogen, which favours a faster chlorination reaction leading to (**64**) and, finally, to 2-iminohydantoins (**66**). Conversely, alkyl groups generate a higher electron density at the amide group, which causes slower rates of chlorination, and allow a competitive reaction of  $\alpha$ -addition to the isocyanide group to give intermediate 2-chlorooxazolines (**68**), which undergo a ring-opening step and a rearrangement leading to 4-iminohydantoins (**70**). This work constitutes the first example of a regioselective cyclisation controlled by the nature of the substituent on the amide group [21].

Likewise, the coupling of isocyanacetamides (**59**) and arylsulfonyl thiocyanates (**22**) led to imidazole-2-thiones (**74**) without the need of an additional base (Scheme 13) [22].

Interestingly, Marcaccini et al. also found that, in contrast to secondary isocyanacetamides (**59**), *N,N*-disubstituted isocyanacetamides (**75**) react with aryl sulfonyl chlorides (**60**) to give oxazoles (**80**). In this case, the enolisation of intermediate (**76**) allows the addition of a second aryl sulfonyl molecule (**60**) prior to cyclisation of the enolic tautomer (**79**; Scheme 14) [23].


Zhu and collaborators made a remarkable contribution with the development of novel two- and three-component oxazole synthesis [19, 24] involving aldehydes (**81**), isocyanacetamides (**82**) and amines (**49**; Scheme 15). The same methodology was followed to make oxazole-derived cyclophanes with high atom economy in a one-step procedure (Scheme 16) [25]. Moreover, Zhu introduced a related enantioselective methodology using Lewis acid catalysis with a chiral (salen)Al(III)Cl complex (**87**; Scheme 15), [26] or BINOL-derived organophosphoric acids (**88**; Scheme 15) [27, 28].

The substrate scope was then expanded by the introduction of propargylamines (**89**) to achieve alkenyloxazoles (**90**) with a novel substitution pattern in a reaction mediated by a




**Scheme 10** Synthesis of quinazoline derivatives


**Scheme 11** Synthesis of mesoionic 3-alkyl-2-arylthio-1,3-diazolium-4-olates from isocyanamides



A) Formation of 2-iminohydantoins:





B) Formation of 4-iminohydantoins:




Scheme 12 Regioselective synthesis of iminohydantoins

Scheme 13 Synthesis of imidazole-2-thiones




Scheme 14 Trisubstituted oxazole synthesis by means of isocyanoacetamides



Scheme 15 Zhu's oxazole synthesis from isocyanoacetamides

Scheme 16 Cyclophanes synthesised from aldehydes, double amines and double isocyanides




stoichiometric amount of  $ZnBr_2$  (Scheme 17) [29]. The transformation, wherein the propargylamine (**89**) acts as a vinyl cation synthetic equivalent, involves a domino sequence incorporating a 1,5-hydride shift, trapping of the in situ generated iminium salt (**93**) by the isocyanide (**82**), cyclisation to the corresponding oxazole (**96**), and 1,6-elimination to yield alkenyloxazole (**90**). This strategy has been used to prepare oxazoles incorporating a steroid skeleton, known to be potent P450<sub>17 $\alpha$</sub>  inhibitors.


Furthermore, oxazole derivatives have been exploited as starting materials for post-condensation transformations. For example,  $\alpha$ -ketoamides (**102**) can be produced through acid hydrolysis of 2-acyl oxazoles (**101**; Scheme 18) [30].

The oxazole scaffold can also be used as a diene in post-Ugi inter- or intramolecular Diels–Alder type cycloadditions. The dienophile is introduced, either as part of one of the starting materials of the Ugi condensation, usually the amine component, or incorporated as external reagent.






Scheme 17 Propargylamines in oxazole synthesis from isocyanoacetamides

Scheme 18 Synthesis of  $\alpha$ -ketoamides from isocyanoacetamides

This approach has been used for synthesis of different heterocycles and other complex compounds, such as pyrrolopyridines (**103**) [31], hexasubstituted benzenes (**104**) [32], tetrahydroquinolines (**105**) [33], phenantrolines (**106**)


[34], polycyclic natural product-like scaffolds (**107**) [35], furoquinolines (**108**) [36], tetrahydrofuropyridines (**109**) [37], and naphthyridines (**110**) [38] (Fig. 1).



**Fig. 1** Heterocyclic scaffolds from oxazole-Diels-Alder reaction pathway

Remarkably, Zhu and Fayol developed the synthesis of an isocyanoacetamide containing a dienophile motif (**113**) that made possible a post-Ugi Diels-Alder cycloaddition resulting in 6-azaindolines (**120**; Scheme 19) [39].

Aiming to access isocyanoacetamides of different natures, Marcaccini and collaborators have previously designed an alternative reaction pathway to synthesise novel isocyanoamides (**123**) through the dehydration of *N*-formamide Ugi adducts (**122**; Scheme 20) [40]. This work is a good




**Scheme 19** 6-Azaindoline synthesis by isocyanoacetamide chemistry


example of the potential of the post-condensation transformation of Ugi adducts. Furthermore, the prepared isocyanoacetamides showed interesting biological properties, for example, cyclic isocyanoacetamides **124** showed good in vitro antimicrobial activity towards *C. albicans* and other microbial agents [41, 42]. More importantly, they are suitable substrates for further use in different isocyanide-based reactions.

In fact, this ingenious strategy moved Marcaccini to explore more exotic isocyanoacetamides for heterocyclic construction. For instance, the condensation between novel isocyanide (**127**) and arylsulfenyl thiocyanates (**22**) furnished 1,3-diazaspiro-2-thiones (**128**), which can exist in three different tautomeric forms (Scheme 21) [43].


In addition, 2,3-disubstituted spiroimidazolones (**133–135**) were formed by *n*-butyllithium treatment of



**Scheme 20** Isocyanide synthesis through post-condensation reaction of Ugi adduct



**Scheme 21** 1,3-Diazaspiro-2-thiones construction through novel isocyanoamides



**Scheme 22** Synthesis of 2,3-disubstituted spiroimidazolones



isocyanoacetamides (**131**) and subsequent trapping of the resulting carbanion with  $\text{NH}_4\text{Cl}$  (**132**) or aldehydes (**81**; Scheme 22) [43, 44].

Bischoff et al. expanded the scope of attainable 2,3-substituted spiroimidazolones by trapping the carbanion intermediate with electrophiles other than aldehydes, such as ketones, amides, or disulphides (**137**) to give 3-substituted spiroimidazolones (**138**). Furthermore, treatment of carbanion with NBS (**39**) introduced a bromine atom resulting in brominated imidazolones (**139**), which were then further subjected to Suzuki or Sonagashira couplings, broadening the diversity of spiroimidazolone products (**141**, **143**;

Scheme 23) [45]. Bischoff also employed a palladium and copper catalysed C–H functionalisation of imidazolones to synthesise analogues of fatty acid synthetase (FAS) inhibitors (**141x**) [46].

An analogous strategy was performed by Pirali's group in the construction of a key intermediate (**152**) in the synthesis of CGRP receptor antagonists. In this sequence, after the Ugi adduct dehydration, the corresponding isocyanoacetamide (**146**) was made to react with a suitable benzyne (**148**) to give intermediate (**149**), which affords the final imidazolone (**152**) through a cyclisation/hydrolysis sequence (Scheme 24) [47].

**Scheme 23** Diversity expansion of 2,3-substituted spiroimidazolones



**Scheme 24** Key intermediate of CGRP antagonists' synthesis by means of isocyanoamide from an Ugi adduct

Zhu and Pirali chose the same synthetic procedure to afford  $\alpha,\alpha$ -disubstituted  $\alpha$ -isocyanoacetamides (**156**), which reacted in a three component reaction to give 5-iminooxazolines (**158**). Furthermore, the reaction between aldehydes (**81**), suitable amino alcohols (**157**) and isocyanoacetamides (**156**), followed by saponification and acid cyclisation resulted in different sized macrocyclodepsipeptides (**159**; Scheme 25) [48].

It is worthwhile to mention here some further studies that helped develop Marcaccini's proposal. For example, Savic *et. al.* proposed the use of DBU as a base and a  $\text{PPh}_3/\text{CBr}_4$  system as a dehydrating agent to transform *N*-formamide Ugi adducts (**160**) into 2-unsubstituted imidazolones (**161**; Scheme 26) [49]. Additionally, Meier *et al.* have recently revisited several isocyanide syntheses and have proposed *p*-TsCl as a cheaper and greener dehydrating agent [50].

## 2,2'-Diethoxy-isocyanoethane

Along with the use of isocyanoacetate (**1**) and isocyanoacetamides (**59**) in the construction of heterocyclic scaffolds, Marcaccini synthesised for the first time 2,2-diethoxy-1-isocyanoethane (**164**) through dehydration of the corresponding *N*-formamide. He then reacted this novel isocyanide with sulphur electrophiles (**60**) or chloramine T (**50**), followed by an intramolecular cyclisation in acidic medium,

to synthesise substituted imidazole cores (**167** and **169**) (Scheme 27) [51, 52].


Marcaccini also reacted 2,2-diethoxy-1-isocyanoethane (**164**) in an Ugi four-component condensation (U-4CC) together with cycloketones (**130**), amine hydrochlorides (**171**) and potassium thiocyanate or selenocyanate (**170**), to obtain spiroimidazo[1,5-*a*]imidazole-5-thiones (**173**) in the presence of acetic acid (Scheme 28) [53].

2,2-Dialkoxy-1-isocyanomethanes (**174**) have facilitated the access to different heterocyclic skeletons. They were even categorised as universal isocyanides for heterocyclic synthesis by Dömling in 2014, who synthesised several Ugi tetrazole derivatives (**178**, **179**, **180**) using **174** in an Ugi-tetrazole reaction followed by acid mediated cyclisation of the Ugi adduct (**177**; Scheme 29) [54].

Kazmaier synthesised thiazoles (**183**) following a thio-Ugi/cyclisation sequence, in which the ring closing stage was similar to Marcaccini's works (Scheme 30) [55], demonstrating the synthetic potential of this isocyanide (**174**).


Moreover, chiral imidazoles (**190**) have been diastereoselectively obtained by Nenajdenko using a similar strategy (Scheme 31) [56]. This reaction readily provides a key intermediate in the synthesis of orally bioactive HIV-1 protease inhibitor SB203386 (190x).


Hulme and Gunawan also took advantage of diethoxyisocyanoethane (**174**) synthetic features to achieve an Ugi/*N*-acyliminium ion cyclisation cascade to afford tricyclic





**Scheme 25** Macrocyclodepsipeptides construction employing isocyanoacetamides

**Scheme 26** Ugi adduct *N*-formamide dehydration and cyclisation with novel base-dehydrating agent system







Scheme 27 2,2-Diethoxyisocyanohydrin in heterocyclic synthesis

Scheme 28 Synthesis of spiroimidazo[1,5-*a*]imidazole-5-thiones

Scheme 29 Synthesis of diverse heterocyclic scaffolds employed 2,2-diethoxyisocyanohydrin

**Scheme 30** Kazmaier's synthesis of thiazoles**Scheme 31** Nenajdenko's synthesis of chiral imidazoles**Scheme 32** Synthesis of brevinamides M-N and fumiquinazolines A-C via isocyanide chemistry

system (**198**), characteristic of marine alkaloids brevianamides M–N and fumiquinazolines A–C (Scheme 32) [57].

### Alkyl (Z)-3-(dimethylamino)-2-isocyanoacrylates

Alkyl (Z)-3-dimethylamino-2-isocyanoacrylates (**199**), or Schöllkopf's isocyanides, were synthesised by Meerwein in 1961 [58]. These are highly versatile isocyanides for heterocyclic synthesis that enable post-condensation transformations of MCR adducts, due to concurrence of isocyanide, alkene and ester functionalities together with a Michael acceptor and a dimethylamino leaving group [59].

Schöllkopf, popularised their use [2] and reported their first application in heterocyclic synthesis [60], employing (**199**) and H<sub>2</sub>S in the preparation of thiazole (**204**; Scheme 33). The mechanism of this reaction involves the formation of an intermediate methanethioamide (**201**) and the subsequent intramolecular Michael cyclisation with elimination of dimethylamine.

Schöllkopf also reported the formation of imidazole rings from isocyanide (**199**) and alkyl or acyl halides [60, 61]. However, Marcaccini et al. revisited this synthesis and showed that the structures proposed by Schöllkopf were incorrect. Instead, substituted oxazoles (**207**) were shown to be obtained by reacting alkyl (Z)-3-dimethylamino-2-isocyanoacrylates (**205**) with acyl chlorides (**100**), as evidenced by their physical and spectral properties [62] (Scheme 34 A). Moreover, Marcaccini also reported the formation of oxazoles (**209**) when electron-deficient arylsulfonyl chlorides (**60**) were used as electrophiles. The mechanism of


this reaction involves the attack of the isocyanide (**205**) on the electrophile and a subsequent ring closing step concerning not the dimethylamino group, but instead the ester group (Scheme 34) [63]. As far as we know, Marcaccini et al. were the only research group that reported oxazole synthesis from Schöllkopf's isocyanide.

Marcaccini et al. also attempted to obtain the imidazole core from Schöllkopf's isocyanide. Thus, in contrast with previous results, reaction of isocyanide **205** with relatively electron-rich arylsulfonyl chlorides (**60**) gave access to the desired imidazole ring, which was then trapped by a second isocyanide molecule and further evolved into the final imidazoloxazolones (**215**; Scheme 35) [64].


The imidazole scaffold was also achieved by Helal and Lucas, by coupling Schöllkopf's isocyanide (**199**) with primary amines (**114**). The reaction takes place with hindered or unhindered amines and proceeds with high regioselectivity and good functional group tolerance, achieving the final products (**218**) in moderate to good yields (Scheme 36) [65].

Moreover, pyrazine core could be built through alkyl (Z)-3-dimethylamino-2-isocyanoacrylate chemistry. For example, Bienaymé and Bouzid achieved pyrazines (**223**) in one pot through an Ugi-tetrazole reaction from Schöllkopf's isocyanide (**220**). In this process, an intramolecular Michael addition took place on the Ugi adduct (**221**) with a release of Me<sub>2</sub>NH to get the final product (Scheme 37) [66].

Later, Illgen et al. [67] developed an Ugi reaction with Schöllkopf's isocyanide (**220**) for pyrazine construction. In this case, the reaction pathway did not involve a Mumm rearrangement of the primary adduct (**225**), but instead an




Scheme 33 Thiazole construction from alkyl-(Z)-2-dimethylamino-2-isocyanoacrylate




Scheme 34 Synthesis of substituted oxazoles







Scheme 35 Synthesis of imidazolyloxazolones




Scheme 36 Imidazole synthesis from Schöllkopf's isocyanide




Scheme 37 Synthesis of bicyclic pyrazines



Scheme 38 Pyrazine construction employing Schöllkopf's isocyanide



**Scheme 39** Ugi post-condensation reaction for thiazole synthesis from Schöllkopf's isocyanide



**Scheme 40** Synthesis of 2,4 substituted thiazoles by isocyanide chemistry catalysed by PPL

intramolecular Michael addition led to the final pyrazine (**227**; Scheme 38).

Schöllkopf's isocyanide (**220**) also readily gives Ugi and Passerini condensations with thiocarboxylic acids (**181**) [59, 68] and the resulting adducts (**228**) have been extensively employed to synthesise thiazoles by the post-condensation transformations. Hence, Dömling reported a thiazole (**231**) synthesis by the cyclisation of the mercaptoimine tautomer (**229**) of the Ugi adduct (**228**; Scheme 39) [69, 70]. This reaction has also been performed on solid support in a combinatorial fashion for the construction of chemical libraries [71].

Recently, an enzyme-catalysed version of this synthesis has been reported by Zhang, (Scheme 40) [72]. Porcine pancreatic lipase (PPL) showed good catalytic activity, allowing the reaction to take place in mild conditions and improving yields of thiazole (**234**) in a reduced reaction time.

Moreover, 1-thiazole-2-yl-methyl-azetidin-2-ones (**241**) were obtained by means of a U-4CC between Schöllkopf's isocyanide (**220**), aldehydes (**185**) and  $\beta$ -aminothiocarboxylic acids (**235**) that facilitated the formation of a  $\beta$ -lactam ring through a 7-membered intermediate (**238**). Then, the

thiazole is formed via Michael addition and elimination of  $\text{Me}_2\text{NH}$ , as previously described (Scheme 41) [73].


Dömling applied an Ugi reaction with Schöllkopf's isocyanide (**220**) to construct analogues of Bacillamide C (**247**), a microbial natural product with algicide and antibacterial bioactivity. The synthetic pathway is described in Scheme 42 [74].

Additionally, disubstituted thiazoles (**251**) have also been synthesised by a Passerini three-component condensation (P-3CC) of oxo-compounds (**153**), thiocarboxylic acids (**181**) and methyl (*Z*)-3-dimethylamino-2-isocyanoacrylate (**220**) under  $\text{BF}_3 \cdot \text{OEt}_2$  Lewis acid catalysis, (Scheme 43) [75].


This strategy was used as a key step in the total synthesis of antimetabolic tubulysin analogues (**254**), as shown in Scheme 44 [76, 77].

## 2-Isocyanothioanisole

Aiming to obtain 2-functionalised benzothiazoles, Marcaccini et al. synthesised a novel isocyanide, 2-isocyanothioanisole (**257**). This compound reacts with



Scheme 41 Synthesis of 1-thiazole-2-yl-methyl-azetidin-2-ones



Scheme 42 Synthesis of Bacillamide C analogues

different electrophiles (**258**) to form intermediates (**259**), which cyclise to yield 5-membered heterocycles (**260**; Scheme 45) [78].


Recently, the chemistry of 2-isocyanothioanisole (**257**) has reached popularity due to its implication in benzothiazole synthesis by means of radical or photochemical approaches.

Wu et al. described the first example of an imidoyl radical coupling with sulphur atom on 2-isocyanoaryl thioethers (**261**). This radical cyclisation can be triggered by a broad scope of radical precursors, such as phosphorus oxides or alkyl radical precursors, yielding 2-substituted


benzothiazoles (**264**) in good yields and broad functional group tolerance (Scheme 46) [79].

Similarly, Liu's group synthesised 2-borylated benzothiazoles (**266**) through radical borylative cyclisation of 2-isocyanothioanisole (**261**) with an *N*-heterocyclic carbene borane (**265**) using AIBN as radical precursor (Scheme 47) [80].

An interesting and more atom-economical alternative was also conceived by Wu and collaborators. This radical cascade employs di-*tert*-butyl peroxide (DTBP) as radical precursor and achieves an intramolecular S to C transfer of the R<sup>2</sup> group, which produces the final 2-substituted benzothiazoles (**264**; Scheme 48) [81].







Scheme 43 Passerini condensation to thiazole construction




Scheme 44 Total synthesis of tubulyisin

Scheme 45 2-Functionalised benzothiazoles from 2-isocyanatoanisole



**Scheme 46** Radical synthesis of benzothiazoles**Scheme 47** Benzothiazole synthesis employing a radical borylative cyclisation**Scheme 48** Radical synthesis of benzothiazoles by reinstallation of the alkyl substituent

Blue LED

**Scheme 49** Photochemical synthesis of benzothiazoles

Radical couplings of 2-isocyanoaryl thioethers (**271**), which produce 2-substituted benzothiazoles (**274**), have also been carried out in photochemical conditions. For example, Yuan et al. synthesised benzothiazoles with fluorine containing motifs in position two (**274**) using visible light, an Ir<sup>4+</sup>/Ir<sup>3+</sup> photocatalyst and Na<sub>2</sub>SO<sub>3</sub> as a reductant (Scheme 49, conditions A) [82]. In a similar way, blue light-mediated fluoroalkylation to obtain substituted benzothiazoles (**274**) was performed with fluoroalkyl iodides using tetramethylethane-1,2-diamine (TMEDA) as electron donor to promote radical coupling with 2-isocyanothioanisole (**271**; Scheme 49, conditions B) [83].

Recently, 4CzIPN (**277**) has gained attention as a photocatalyst for irradiation with blue light to form benzothiazoles from 2-isocyanoaryl thioethers (**271**). Wu et al. have designed a metal-free, oxidant-free protocol for the synthesis of 2-substituted benzothiazoles (**276**) based on a photocatalysed reaction between (**271**) and cyclic or acyclic ethers (**275**) through a SET pathway (Scheme 50) [84].


Lastly, Liu and collaborators have developed a phosphorus radical cascade employing their catalyst 4CzIPN-*t*Bu, a modification of 4CzIPN (**277**) in which *t*-Bu groups have been introduced on positions 3 and 6 of each carbazole motif. They used this catalyst to perform a PECT cycle to

obtain 2-phosphorylated benzothiazoles from 2-isocyanoaryl thioethers (**271**), a reaction that was previously performed without photochemical activation [79]. Notably, they were also able to synthesise diverse heterocycles, such as phenanthridines or quinolines using different isocyanides [85].

## Intramolecular Ugi and Passerini reactions


Intramolecular Ugi reactions are possible when one of the starting materials contains two of the functional groups involved in the reaction. This strategy has been employed by numerous groups to increase the scaffold diversity of the classical Ugi reaction. Marcaccini developed diverse original intramolecular Ugi and Passerini reactions using bifunctional starting materials. In this way, he was able to prepare libraries of different privileged structures with relevance in the field of medicinal chemistry and chemical biology. These developments have had an important influence in the further contributions of other research groups [86, 87].


The use of oxocarboxylic acids (**278**) as bifunctional starting material in IMCR was developed by Short [88] and Harriman [89] for the preparation of five-, six-, seven- and eight-membered lactams (**279**) through Ugi four-centre



**Scheme 50** Photosynthesis of 2-benzothiazoles from 2-isocyanothioanisole and ethers

**Scheme 51** First examples of the use of oxoacids in the Ugi reaction



**Scheme 52** Ugi reaction of 5-oxo-3-thiacarboxylic acids

3-component reactions (U-4C-3CR; Scheme 51). Similarly, Ugi prepared diverse  $\gamma$ -lactams by using levulinic, 3-benzoylpropionic and phthalaldehydic acids. When amino esters are used as the amine component, it is possible to obtain 1,4-diazabicyclo[4.3.0]nonane-3,5,9-triones (**280**) [90].


Stefano Marcaccini described for the first time the Ugi four-centre three-component reaction (U-4C-3CR) between 5-oxo-3-thiacarboxylic acids (**281**), benzylamines (**114**) and cyclohexyl isocyanide (**282**). The reaction takes place in refluxing methanol giving 5-oxothiomorpholine-3-carboxamides (**283**) with good yields and high diastereoselectivities (Scheme 52). The major diastereoisomer has been assigned *trans* configuration by NOESY experiments of the bicyclic compounds (**284**) [91].

This strategy was extended to U-4C-3CR of 6-oxo-4-thiacarboxylic acids (**285**) to form biologically relevant hexahydro-1,4-thiazepin-5-ones (**286**) and 1,4-benzothiazepin-5-ones (**289**). This condensation takes place in some cases with high stereoselectivity (Scheme 53) [92]. Furthermore, chemoselective reduction with  $LiAlH_4/AlCl_3$  of the cyclic carbonyl group gives bicyclic 1,4-thiazepine-3-carboxamides (**287**).


Ivachtchenko used a similar protocol to synthesise 3-oxo-1,4-thiazepine-5-carboxamides (**294**) [93], 5-oxo-1,4-oxazepine-3-carboxamides (**296**) [94] and 5-oxo-1,4-thiazepine-3-carboxamides (**298**) [93] fused to diverse heterocycles from the oxoacids (**293**), (**295**) and (**297**), respectively (Scheme 54, B–D). He also obtained nine-membered heterocyclic scaffolds (**300**) by the reaction of the aromatic aldehyde-acid (**299**; Scheme 54, E) [95]. As well, Zhang et al. have previously used different oxoacids, including 2-(2-formylphenoxy)acetic acid (**291**) to prepare diverse lactams and oxazepines, such as **292** (Scheme 54, A) [96].

On the other hand, the reaction of similar ketoacids (**301**) with isocyanides (**31**) and *Boc*- or *Cbz*-protected hydrazine (**302**) was used by Krasavin for the synthesis of *N*-aminolactams (**303**), which are proline-like  $\beta$ -turn secondary structure mimics (Scheme 55) [97].


Marcaccini performed an intramolecular Passerini condensation between oxothiocarboxylic acid (**304**) and isocyanides (**31**) under tertiary amine catalysis. This reaction affords seven-membered lactones (**306**) that suffer spontaneous intramolecular nucleophilic attack of the amide NH to give unexpected tetracyclic 1,4-benzothioxepin orthoamides (**307**). The observed stereochemistry of the

**Scheme 53** Intramolecular Ugi condensation of 6-oxo-4-thiacarboxylic acids

**Scheme 54** Synthesis of oxazepine and thiazepine-fused carbocycles and heterocycles




**Scheme 55** Synthesis of *N*-amino lactams



major product is due to an axial attack of the isocyanide to yield a *cis*-fused  $\alpha$ -adduct (**305**), which, evolves to give the *trans*-fused Passerini adduct (**306**; Scheme 56) [92]. This is an appealing result as the *N*-amidoalkoxycarbonyl


moiety (orthoamide) was considered an unstable intermediate in the reaction of an ester with an amide anion or an imide with an alkoxy anion. At the time, the only known







**Scheme 56** Intramolecular Passerini condensation to prepare tetracyclic 1,4-benzothioxepin orthoamides

**Scheme 57** Synthesis of lactones by intramolecular Passerini reaction



**Scheme 58** Synthesis of isoindolinone-carboxamide derivatives



**Scheme 59** Obtention of isochromenone Ugi primary adduct and their transformation

stable compounds having this structural feature were peptide ergot alkaloids (ergopeptines).

In contrast, Orru and Ruijter found that when the ketoacid (**308**) or other  $\gamma$ - and  $\delta$ -keto acids are used in a Passerini three-centre two-component reaction (P-3C-2CR), the corresponding *trans*-fused lactone (**310**) is obtained with good diastereoselectivity. Interestingly, these Passerini adducts (**310**) are readily rearranged to less strained *cis*-fused  $\alpha$ -hydroxyimides (**311**) under acidic conditions (Scheme 57) [98].

As with other oxoacids, 2-formylbenzoic acid (**312**) was reported to suffer U-4C-3CRs with amines (**114**), and isocyanides (**31**) to give 2-isoindolinone-7-carboxamide analogues (**313**; Scheme 58) [96, 99].


Remarkably, Marcaccini developed an interrupted Ugi condensation from 2-formylbenzoic acid (**312**), scarcely basic amines (**114**) and isocyanides (**31**). In this reaction, as in the regular Ugi condensation, the nitrilium intermediate is intramolecularly attacked by the carboxylic acid to give the usual primary adduct (**314**). However, in this case, this

intermediate does not suffer a Mumm rearrangement, but is instead stabilised by a tautomerisation leading to isochromenone enediamine (**315**). In this process, the precise control of the reaction conditions (solvent, time, and temperature) is essential to obtain the products (**315**) satisfactorily. In some cases, isocoumarins (**315**) in the presence of a catalytic amount of acid suffer a rearrangement to give isoindolines (**313**) in quantitative yield (Scheme 59) [100].


This reaction was later reported by Ramazani to be catalysed by silica nanoparticles in solvent-free conditions, though the amine component was limited in this case to dibenzylamine [101]. The same protocol was also used to synthesise isochromenone-functionalised mesoporous silica hollow spheres from 2-formylbenzoic acid (**312**), 2,6-dimethylphenyl isocyanide and amine-containing silica spheres [102].

Marcaccini also reported the reaction of isocoumarin enediamines (**317**) with amines (**114**), which promote a ring cleavage that gives new phenylglycine derivatives (**318**) in almost quantitative yields (Scheme 60). This process allows four diversity elements to be introduced through an U-4C-3CR. The reaction was performed in solventless conditions using an excess of the amine to prevent the decomposition of labile isocoumarins (**317**) [103].


An enantioselective version of this process has been described by Zhu using an octahydro (*R*)-BINOL-derived chiral phosphoric acid (**323**). The reaction of 2-formylbenzoic acids (**316**), isocyanides (**31**), and aromatic amines (**49**) in the presence of a catalytic amount of chiral phosphoric acid (**323**) affords the *S* isoindoline (**322**) with good enantioselectivity. The authors justify the observed enantioselectivity by a dynamic kinetic resolution process in which



**Scheme 60** Synthesis of phenylglycine derivatives from primary Ugi adducts



**Scheme 61** Enantioselective intramolecular Ugi reaction

**Scheme 62** Synthesis of pyrazino[2,1-*a*]isindolediones

there is an imine-enamine tautomerisation equilibrium, much faster than acid-catalysed Mumm rearrangement (Scheme 61) [104].

When the propargyl amine (**324**) is used in the Ugi reaction of 2-formylbenzoic acids (**316**) and isocyanides (**31**), it is possible to obtain the pyrazino[2,1-*a*]isindolediones (**327**) by 6-*exo-dig* intramolecular hydroamination on the Ugi adducts (**325**), followed by a 1,3-H shift (Scheme 62) [105].


The reaction of 2-formylbenzoic acid (**312**) with isocyanides (**31**) and amino alcohols or bis-secondary amines (**328**) in methanol, under microwave irradiation, at 60 °C permits the synthesis of eight and nine-membered lactones or lactams (**329**), probably through an isochromenone enediamine. When *L*-prolinol is used as the amine component, diastereomeric lactones (**330**) are obtained in a 1.5:1 ratio (Scheme 63) [106].

A combination of a P-3C-2CR of 2-formylbenzoic acid (**312**) and isocyanides (**31**) and a subsequent aldol condensation with arylglyoxals (**333**) was reported by Jiang and Tu. The Passerini reaction takes place in methanol to give


an intermediate enamine (**332**). This reacts with arylglyoxal (**333**) through an aldol condensation to give hydroxyaldehyde (**334**), which then undergoes an intramolecular nucleophilic addition and a ring-opening process to give the isocoumarins (**336**; Scheme 64) [107].

The authors argue that arylglyoxals (**333**) are mainly in their hydrate form in methanol and therefore are not prone to participate in an intermolecular P-3CC. On the other hand, when the reaction is carried out in dioxane, arylglyoxals exist mostly in the aldehyde form and a P-3CC takes place swiftly to give (**337**), which then evolves through an intramolecular aldol reaction to yield isomeric isocoumarin (**339**; Scheme 65) [107].

A divergent preparation of isocoumarins (**343**) and thiophthalides (**345**) was described by El-Kaïm. Oxoacid (**316**) reacts with thiols (**35**) in the presence of *p*-toluenesulfonic acid catalyst and magnesium sulphate to give 3-sulfanylphthalides (**340**), which suffer an insertion reaction of isocyanide (**31**) mediated by titanium tetrachloride to give the isocoumarins (**343**). When *tert*-butyl thiol (**35**,  $\text{R}^2 = \text{tBu}$ ) is used, thiophthalides (**345**) are obtained by a formal

**Scheme 63** Synthesis of eight and nine-membered lactones and lactams

**Scheme 64** Synthesis of isocumarins by tandem P-3C-2CR / aldol reaction




**Scheme 65** Synthesis of isocumarins by P-3CC followed by aldol reaction

thio-Passerini reaction. The deprotection of the *tert*-butyl group in intermediate (**342**) leads to the thiophthalides (**345**) by a 1,5-Mumm rearrangement. The process is also possible in one pot (Scheme 66) [108, 109].


The experience with isocyanoacetates (**1**) in the synthesis of heterocycles, as mentioned above, surely encouraged Marcaccini to use isocyanoacetic acid in Ugi-type reactions. For example, an intramolecular Ugi reaction between ketones (**153**), amine hydrochlorides (**171**) and potassium isocyanoacetate (**346**) affords the Ugi primary adducts (**347**), which suffer the attack of a second amine molecule to give the unexpected dipeptide derivatives (**348**) in place of the expected ketopiperazines (**349**) product of the Mumm rearrangement (Scheme 67) [110].

Zhu et al. found that this reaction could be extended to the use of secondary amines and aldehydes or ketones if toluene is used as a solvent [111]. They also found that the presence

of ammonium chloride as an additive is crucial for the success of the reaction. Conversely, the reaction of aldehydes or cyclic ketones (**153**) with dimethylamine hydrochloride (**351**) and potassium 2-isocyano-2-aryllacetates (**350**) gave the *N*-acyl imino amides (**355**) as the predominant product (Scheme 68). Subsequent treatment with aqueous acid produces amide (**356**) and ketoamide (**357**). According to the mechanism proposed by the authors, when an aromatic ring is attached to the  $\alpha$ -position of isocyanide (**350**), the oxazolone (**352**) exists in the predominant enolic form (**353**) and suffers a 1,6-elimination of dimethylamine to give (**354**). The ensuing nucleophilic attack of dimethylamine results in the ring-opening to yield *N*-acyl imino amide (**355**), which gives by hydrolysis the amide (**356**) and ketoamide (**357**; Scheme 68). In this case, the reaction takes place in toluene at room temperature without ammonium chloride and only 1.2 equivalents of amine are used [112].



**Scheme 66** Reaction of formylbenzoic acids, thiols and isocyanides




**Scheme 67** Synthesis of dipeptides from potassium isocynoacetates

## Post-condensation transformations of Passerini and Ugi adducts

Marcaccini performed extensive work on the transformation of Passerini and Ugi adducts. His work makes clear that a well-planned sequence of classical Ugi or Passerini isocyanide multicomponent reactions and post-condensation transformations constitutes an extremely powerful synthetic methodology for the preparation of structurally diverse complex molecules, such as heterocyclic compounds with elaborate substitution patterns, constrained

peptides, peptide mimetics, and pseudopeptides. The presence of complementary reactive groups in the multicomponent adducts may facilitate post-condensation transformations that depend on the nature and position of these groups. These reactive groups can easily be incorporated into the product as part of some of the MCR reagents. Generally, protecting group strategies are not necessary, as IMCRs are tolerant to a great diversity of functional groups. Interestingly, some functionalities created in the IMCR can be used in further transformations, meaning that only one extra functional group must be present in one of the starting reagents. Thus, two acidic positions in



**Scheme 68** Transformation of oxocompounds to amides


the Ugi and Passerini adducts –the NH amide group and the peptidyl CH position — can undergo deprotonation, generating new nucleophilic centres able to react intramolecularly with other functional groups.

### Enolisation of the peptidyl hydrogen

In some cases, the peptidyl hydrogen on the asymmetric carbon resulting from a Passerini or an Ugi reaction can be abstracted with a base to generate a nucleophilic carbanion

able to react intramolecularly with electrophilic groups in the adduct. However, in most cases, this hydrogen is not acidic enough, despite being adjacent to an electron-withdrawing amide. Marcaccini succeeded in increasing the acidity of the peptide hydrogen by introducing functionalities capable of stabilizing the resulting carbanion. For example, the use of cinnamaldehyde (358) as one of the components in a four-component Ugi condensation affords an adduct (360) that it is easily deprotonated with moderate bases to give a highly delocalised anion (361), containing several

**Scheme 69** Synthesis of  $\beta$ -lactams from Ugi adducts



reactivity centres. When a complementary electrophilic centre is introduced with one of the components of the U-4CC, such as chloroacetic acid (**359**), the adduct (**360**) can be easily cyclised with KOH in methanol to a  $\beta$ -lactam (**362**; Scheme 69) [113]. This is a rare example of  $\beta$ -lactam ring formation via a C<sub>3</sub>-C<sub>4</sub> bond.

Remarkably, a similar strategy for the synthesis of 4-phosphono- $\beta$ -lactams (**365**) was later reported by Stevens et al. (Scheme 70) [114, 115].


It is interesting that in both Marcaccini's and Stevens' syntheses, cyclisation occurs by attack of the  $\alpha$ -carbon rather than the  $\gamma$ -carbon. This seems counterintuitive, as when there is an intramolecular ring-closure competition between a four- and a six-membered ring, the latter is usually

preferentially formed. However, in this case, the four-membered ring is selectively formed. Theoretical calculations carried out by Van Speybroeck, Stevens et al. suggest that the four-membered ring preference is due to a geometrically strained S<sub>N</sub>2-like transition state for the six-membered ring formation [116].


Another very similar approach to  $\beta$ -lactams reported by González-Muñiz starts from amino esters (**366**) that are acylated with chloroacetyl chloride (**367**) to give (**368**) and then cyclised in a basic medium to the corresponding  $\beta$ -lactams (**369**; Scheme 71) [117, 118].

Interestingly, the  $\beta$ -lactams obtained with Marcaccini's procedure are stable when the amide nitrogen is substituted with a relatively electron-rich aromatic group resulting from


**Scheme 70** Synthesis of 4-phosphono- $\beta$ -lactams reported by Stevens



**Scheme 71** Synthesis of  $\beta$ -lactams from natural amino acids




**Scheme 72** Synthesis of succinimides from Ugi adducts




non-substituted anilines or anilines containing electron-donating groups (**49a**; Scheme 69). On the other hand, the use of anilines with strong electron-withdrawing substituents (**49b**) results, after basic treatment of the adducts (**360b**), in  $\beta$ -lactams containing a good leaving nitrogen group (**362b**) that cannot be isolated and are spontaneously transformed to succinimides (**370**) in the reaction conditions (Scheme 72). Thus, the same synthetic procedure gives selective access to two important heterocyclic scaffolds depending on the particular reagents used in each case [119, 120].

Years later, Balalaie et al. used this same tactic for the synthesis of  $\beta$ -lactams (**374**) and succinimides (**375**) using the Ugi adducts (**373**) of phenylpropionic acid (**372**). In this case, an aromatic ring is sufficient to stabilise the peptidyl anion, which is generated by  $K_2CO_3$ . These authors were able to isolate the  $\beta$ -lactams (**374**) when the cyclisation is performed in acetonitrile, while the isomerised succinimide products (**375**) are obtained in methanol (Scheme 73) [121]. Very recently, Wang, He and co-workers substituted but-3-ynoic acid derivatives for phenylpropionic acid, obtaining




**Scheme 73** Balalaie's version of Marcaccini's synthesis of  $\beta$ -lactams and succinimides

**Scheme 74** Synthesis of 1,6-dihydro-6-oxopyridines



**Scheme 75** Gómez-Montaña—El Kaïm synthesis of aminopyrrolinones





$\gamma$ -lactams by U-4CC followed by an intramolecular 5-*exo-dig* cyclisation in the presence of  $\text{Cs}_2\text{CO}_3$  [122].

Moreover, Marcaccini reported that the Ugi adducts of cinnamaldehyde (**358**), glyoxylic acids (**376**), amines (**114**) and isocyanides (**31**) can be easily transformed into oxopyridines (**380**) by basic treatment (Scheme 74) [123]. The key intermediate is again the highly stabilised anion (**378**) which, in this case, behaves as a benzylic anion in the intramolecular nucleophilic attack on the carbonyl group.

Here again, a simple aromatic group can also sufficiently stabilise the peptidyl anion, making further cyclisation to heterocyclic products possible. Gómez-Montaño and El Kaim were able to cyclise in this way the cyanoacetic acid-derived Ugi adducts (**382**), previously prepared by Marcaccini [124–127], to give aminopyrrolinone derivatives (**383**; Scheme 75) [128].

Ivachtchenko et al. used the acidity of the Ugi peptidyl hydrogen to promote aromatic nucleophilic reactions leading to isoindole derivatives (**385**). The reaction takes place with triethyl amine in DMF at high temperatures. In this case the anion is stabilised either by a phenylvinyl group, as in the previous Marcaccini's reactions, or by



electron-deficient aromatics introduced as the aldehyde component in the U-4CC (Scheme 76) [129].

Liu et al. used a very similar strategy to prepare 3-(indol-2-yl)isoindolin-1-ones by the cyclisation of the adducts of 1*H*-indole-2-carbaldehyde, 2-iodobenzoic acid, amines and isocyanides. In this case the reaction was carried out in DMSO, under microwaves, at 80 °C, using  $\text{Cs}_2\text{CO}_3$  as a base [130].

The possibility of generating a nucleophilic peptidyl anion allows for the synthesis of a variety of scaffolds limited only by the nature of the electrophilic groups present in the Ugi adducts. For example, Miranda synthesised 2,3-dihydropyrroles (**390**) by the cyclisation of peptidyl anions generated from Ugi-allenamide adducts (**388**). The latter in turn are prepared in situ from propargyl amine U-4CC adducts (**387**; Scheme 77) [131]. The utility of this methodology was demonstrated by Vázquez, who used it in conjunction with catalytic hydrogenation of pyrroline products to synthesise a library of nicotine analogues (**390h**) [132].

The same strategy was recently applied by Balalaie et al., who cyclised the Ugi adducts (**392**) of allenic acids (**391**)

**Scheme 76** Ivachtchenko's synthesis of 3-oxoisoindoline-1-carboxamides



**Scheme 77** Miranda's synthesis of 2,3-dihydropyrroles

to obtain pyrrolidin-5-one-2-carboxamides (**393**), this time through a 5-*exo*-dig approach (Scheme 78) [133].

A similar method, which starts from 3-chloropropionic acid (**395**) was further developed by Vázquez et al. for the synthesis of cotinine and iso-cotinine analogues (**398**). Here, the direct displacement of the chlorine atom by the attack of the peptidyl anion could explain the formation of the products, but instead evidence supports a two-step base-mediated elimination/Michael addition mechanism (Scheme 79) [134]. A very similar approach to the synthesis of indolyl-substituted lactams by the cyclisation of Ugi-adducts in basic conditions has been also published by Shire et al. [135].



An interesting application of the previous reaction was developed by Yang et al. for the diastereoselective synthesis of chromeno[3,4-*c*]pyrrole-3,4-diones (**402**). The carboxylic

component of the U-4CC is the  $\alpha,\beta$ -unsaturated chromene-3-carboxylic acid (**400**) and the resulting adduct (**401**) spontaneously suffers an intramolecular Michael addition to the expected chromenopyrrole (**402**; Scheme 80) [136].


The intermolecular addition of Ugi peptidyl carbon to Michael acceptors has also been achieved by Abderrahim and El Kaïm in a two-step synthesis of pyrrolines (**407**). The authors suggest that a concerted [3 + 2] mechanism involving the formation of a dipolar derivative of the Ugi adduct (**404**) occurs (Scheme 81) [137]. Analogously, the peptidyl carbon on Passerini adducts has been shown to act as a nucleophile in Michael additions with acrylonitrile. The resulting  $\gamma$ -hydroxynitrile can be then be cyclised under acidic conditions to yield  $\gamma$ -butyrolactones [138].

El Kaïm's group also achieved a simple and general synthesis of  $\beta$ -lactams (**411**) by the addition of diiodomethane

**Scheme 78** Balaia's synthesis of pyrrolidinones




**Scheme 79** Vázquez's synthesis of cotinine analogues




**Scheme 80** Diastereoselective synthesis of chromeno[3,4-*c*]pyrrole-3,4-diones

**Scheme 81** Formal [3+2] cycloaddition of Ugi adducts for the synthesis of pyrrolines



**Scheme 82** Synthesis of  $\beta$ -lactams by  $\text{CH}_2\text{I}_2$  addition to Ugi adducts



(408) to amide dianions (409) obtained by deprotonation of the Ugi adducts (403; Scheme 82) [139]. Calculations suggest that  $\text{CH}_2\text{I}_2$  (408) is first added to the peptidyl carbon and the resulting iodomethane-substituted amide (410) then cyclises to the desired  $\beta$ -lactam (411).


Analogously, propargyl bromide (415) was used as bis-electrophile to trap Ugi amide dianions (414), resulting in the formation of pyrrolidinone enamides (416). This reaction has been combined with a subsequent Pictet–Spengler cyclisation to give benzoindolizine scaffolds (417) present in the heterocyclic core of crispine alkaloids (Scheme 83) [140]

Other strategies to activate the peptidyl carbon of Ugi adducts have been developed by different groups. As part of their research in the elaboration of Ugi adducts by means of transition metal-catalysed reactions, Neuville and Zhu reported the synthesis of 3-substituted 3-benzoxazolyloisoindolinones (422) starting from an Ugi-adduct having two aryl iodide units (420). The process takes place

through a regioselective sequential intramolecular copper-catalysed *O*-arylation and palladium-catalysed *C*-arylation of the adduct peptidyl position (Scheme 84) [141].



A similar strategy was recently developed by Ghandi et al. for the synthesis of spiropyrroloquinoline isoindolinones and aza-isoindolinones (426). In this case, the double cyclisation takes place under metal-free conditions (Scheme 85) [142].

A different double cyclisation was proposed by Van der Eycken for the synthesis of spiroindolinone-isoindolinone derivatives (430) from Ugi adducts (229) in the presence of a palladium catalyst and a base. The authors propose a reaction mechanism involving a first palladium-catalysed Buchwald–Hartwig C–N coupling, followed by a base-promoted addition of the peptidyl carbon to the remaining aryl halide (Scheme 86) [143]. Bromobenzoic acid (428) may also be replaced by propionic acids in the Ugi reaction. In this latter case, spiroindolinone-pyrrolones



Scheme 83 Ugi/propargylation/Pictet–Spengler cyclisation


Scheme 84 Palladium-catalysed intramolecular C-arylation of the Ugi peptidyl carbon



Scheme 85 Ghandi synthesis of spiranes


are formed through a Buchwald–Hartwig/Michael addition sequence [144].

On the other hand, El Kaïm and Miranda achieved a copper-catalysed oxidative activation of the peptidyl position of Ugi adducts (**432**) leading to a double radical coupling to




**Scheme 86** Van der Eycken synthesis of spiranes

**Scheme 87** Radical peptidyl activation leading to spiroindolines



**Scheme 88** Deamidative  $\text{C}(\text{sp}^2)\text{-C}(\text{sp}^3)$  coupling for the synthesis of isoindolinones




efficiently produce complex polycyclic spiroindolines (**437**; Scheme 87) [145].

More recently, Ali and El Kaïm carried out the allylation of Ugi adducts at their peptidyl position using allyl acetate in both the presence and absence of palladium catalysts. This strategy allows the preparation of bis-alkenyl derivatives suitable for subsequent ring-closing metathesis leading to nitrogen heterocycles [146].


An interesting copper-catalysed intramolecular coupling between the peptidyl carbon and aryl iodide was developed by Chauhan et al., providing a straightforward synthesis of isoindolinones (**439**). The coupling takes place

with concomitant loss as isocyanate of the amide moiety originally introduced with the isocyanide component of the U-4CC (Scheme 88) [147]. The mechanism of this reaction is obviously different from the mechanism of Ivachtchenko's approach to isoindolinones, in which no deamidation is produced (Scheme 76) [129]. Van der Eycken proposed a variant of this reaction using 2-chloronicotinic acid-derived Ugi adducts that cyclise in the presence of a base, with no need of metal catalysis [148].


Marcos et al. have recently developed an oxidative C(sp<sup>3</sup>)-H intramolecular imination of hydroxycoumarin enol-Ugi adduct derivatives (**442**) which leads to



Scheme 89 Amide-directed oxidative cyclisation of enol-Ugi derivatives



Scheme 90 Synthesis of 2(3H)-oxazolone 4-carboxamides



**Scheme 91** Synthesis of  $\beta$ - and  $\gamma$ -lactams by the Ding's group

imidazolocoumarins (**447**) [149]. Interestingly, the regioselectivity of the reaction is controlled by the amide group derived from the enol-Ugi isocyanide component, which directs the functionalisation of the adjacent  $C(sp^3)-H$  and then is lost as an isocyanate (Scheme 89).

In hopes of creating easily enolisable Ugi adducts, Marcaccini introduced the use of arylglyoxals (**333**) as the carbonyl component. The presence of an additional carbonyl group at the enolisable position of the adduct would obviously favour the formation of the corresponding enolate. Thus, the U-4CC of arylglyoxals (**333**), anilines (**49**), isocyanides (**31**) and trichloroacetic acid (**448**) led to highly reactive Ugi adducts (**449**) that spontaneously cyclised in the reaction medium to yield oxazolones (**450**; Scheme 90) [150]. Interestingly, Marcaccini introduced here for the first time the trichloromethyl functionality as a convenient leaving group in post-condensation transformations of IMCRs.

The ready enolisation of the Ugi adducts of arylglyoxals (**452**) was used by Ding in a synthesis of  $\beta$ -lactams (**453**) conceptually related to those published by Marcaccini in 1998 (Scheme 91, A) [119, 151]. In this case, the leaving group is a bromine atom from the bromoacetic acid (**451**) component used in the U-4CC. The use of a 3-bromopropionic acid derivative (**454**) permitted, in turn, the synthesis of the corresponding 5-membered lactams (**455**; Scheme 91, B) [152]. Analogously, the related Passerini reaction led to the corresponding  $\gamma$ -lactone [153].



Interestingly, the  $\beta$ -lactams (**457**) obtained from the Ugi adducts of  $\alpha$ -activated amines (**456**), chloroacetic acid (**359**)

and arylglyoxals (**333**), are easily transformed to highly functionalised  $\gamma$ -lactams (**460**) by treatment with simple and low-moisture-sensitive bases. The ring expansion is explained by an anionic rearrangement. Thus, the initial deprotonation of the acidic position in the *N*-substituent of the azetidinone (**457**) triggers a ring opening that yields a new intermediate containing an imine and an enolate, which cyclises to the  $\gamma$ -lactams (**460**; Scheme 92) [154].

Peshkov and Van der Eycken proposed the use of arylglyoxals (**333**) and propiolic acids (**461**), as aldehyde and acid components, respectively, to prepare easily enolisable Ugi adducts also containing a Michael acceptor (**462**). The U-4CC, carried out in methanol at 80 °C, was spontaneously followed by the intramolecular Michael addition of the peptidyl anion to the triple bond to give the corresponding pyrrolones (**463**). A subsequent retro-Claisen fragmentation results in the cleavage of the benzoyl moiety to yield the final pyrrolone-2-carboxamides (**464**; Scheme 93) [155]. Here, in contrast to Balalaie's work cited above [121], a *5-endo-dig* rather than a *4-exo-dig*-carbocyclisation takes place.


In a subsequent work, the same group substituted heterocyclic aldehydes containing basic nitrogen atoms for the glyoxal carbonyl component of the U-4CC, obtaining the expected  $\gamma$ -lactams [156]. Moreover, an analogous Passerini/cycloisomerisation process allowed to conveniently obtain butenolides (**467**) in good yields (Scheme 94) [157].

Taking advantage of the ready enolisation of the Ugi adducts derived from arylglyoxals as well as their feature as doubly functionalised reactants, García-Valverde et al.

Scheme 92 Ring expansion of  $\beta$ -lactams

Scheme 93 Peshkov and Van der Eycken synthesis of pyrrolone-2-carbox-amides

Scheme 94 Peshkov and Van der Eycken synthesis of butenolides



have developed a one-pot synthesis of enantiopure pyrrolpiperazines (**471**) through a diastereoselective multicomponent domino reaction, Ugi/enamine alkylation, using non-protected 1,2-diamines (**469**) together to arylglyoxals (**333**) and 3-bromopropionic acid (**468**). Thus, after the formation of a cyclic diamine, the Ugi reaction takes


place selectively over the aldimine position. The spontaneous alkylation of the cyclic enamine intermediate (**470**) affords the corresponding pyrrolpiperazine (**471**), with the generation of a new stereogenic centre in the last step (Scheme 95) [158].




## Amide NH as nucleophile

The transformations shown above rely on the formation of a highly stabilised anion in the peptidyl position. This occurs when the appropriate substituents are introduced on Passerini or Ugi adducts with the initial reaction components. However, when the aldehyde employed as starting reactant is unable to stabilise the peptidyl anion, or this anion is not


in the correct position to react with other groups in the molecule, a base-induced transformation involving the amide nitrogen may take place. An interesting example reported by Marcaccini is the basic treatment under ultrasonic sonication of the Ugi adduct (**472**) of aromatic aldehydes (**370**) and chloroacetic acid (**359**), which affords 2,5-diketopiperazines (**474**) as a result of an intramolecular *N*-alkylation (Scheme 96) [159].




Scheme 95 Multicomponent domino reaction in the synthesis of pyrrolo-piperazines




Scheme 96 Synthesis of 2,5-diketopiperazines



Scheme 97 Diastereoselective synthesis of 2,5-diketopiperazine-derived steroids



**Scheme 98** Synthesis of homochiral 2,5-diketopiperazines



**Fig. 2** 2,5-Diketopiperazine inhibitors of Apaf-1

Marcaccini described the limitations of this methodology. On one hand, the cyclisation was unsuccessful with aliphatic aldehydes and, on the other, epimerisation of the stereogenic centre occurs in the cyclisation conditions, as evidenced by the incorporation of deuterium at the C3 position when KOD in EtOD was employed. However, when the aldehyde group was directly linked to a steroidal framework (**475**), these drawbacks were not observed and, despite the aliphatic characteristic of the aldehyde employed, the reaction took place in good yield. Moreover, high diastereoselectivity was achieved in the Ugi reaction, possibly induced by


the rigid steroidal frame, and maintained during the cyclisation (Scheme 97) [160].

Interestingly, Banfi et al. described a similar strategy for the synthesis of 2,5-diketopiperazines (**480a** and **480b**) from chiral amino alcohols (**478**) using caesium carbonate instead of potassium hydroxide in the final stage. The cyclisation was carried out on the separated diastereomers of the Ugi adducts (**479a** and **479b**), as under these reaction conditions the epimerisation on the C3 position of the diketopiperazines was not observed (Scheme 98) [161].

Messeguer et al. employed this methodology in the design of analogues of linear *N*-alkylglycine oligomers (**481**), reported as apoptotic inhibitors [162]. These authors created cyclic motifs in these peptoids to introduce geometric constraints that reduce conformational freedom and increase their selectivity. Moreover, to avoid rotamers, they substituted an isostere triazole moiety for the tertiary amine (**482**; Fig. 2). The resulting peptoids were shown to be Apaf-1 inhibitors that decrease the apoptotic phenotype in mitochondrial-mediated models of cellular apoptosis.

They carried out the synthesis of these restricted analogues (**482**) through an Ugi reaction combining chloroacetic acid (**359**) and triazole aldehydes, followed by a base-induced intramolecular cyclisation, which afforded, not only the expected 2,5-diketopiperazine (**482**), but also the corresponding  $\beta$ -lactam (**484**), because of the increased acidity

**Scheme 99** Divergent cyclisation of chloroacetic acid derived Ugi adducts




on the peptidyl position (Scheme 99) [163]. The influence of the substitution on the secondary amide and the triazole core was determinant for the experimental results and was theoretically studied [164, 165]. Interestingly, the peptidomimetics bearing the  $\beta$ -lactam scaffold (**484**) turned out to be more potent apoptotic inhibitors than the diketopiperazine isomers.

Additionally, diketopiperazines fused with other heterocycles have been synthesised by Dehaen et al. using this methodology, combining chloroacetic acid with other functionalised reactants in the U-4CC. Thus, triazolobenzodiazepine diketopiperazines (**489**) were synthesised from 2-azido-benzaldehydes (**485**), propargyl amines (**486**), chloroacetic acid derivatives (**487**) and isocyanides (**31**). The Ugi adduct was cyclised under basic conditions to the diketopiperazine


system bearing azide and alkyne functionalities (**488**), which was then subjected to intramolecular azide-alkyne cycloaddition (IAAC) in refluxing ethanol to give the fused system (**489**; Scheme 100) [166].

Miranda used 2-bromobenzylamines (**491**) and benzoyl-acetaldehyde (**490**) as doubly functionalised reactants along with chloroacetic acid (**359**) in the synthesis of pyrazinoisoquinolines (**493**). The fused systems were obtained through a three-step protocol consisting of an Ugi reaction followed by a basic medium-promoted cyclisation/elimination and a 6-*endo* Heck cyclisation (Scheme 101) [167].


The synthesis of tetrahydropyrazino[1,2-*a*]indole-1,4-diones (**497**) by the Joullié-Ugi reaction of indolenines (**494**), haloacetic acids (**495**) and isocyanides (**31**), followed by an intramolecular *N*-alkylation, was reported by Krasavin



**Scheme 100** Synthesis of polycyclic diketopiperazines reported by Dehaen



**Scheme 101** Synthesis of pyrazinoisoquinolines



**Scheme 102** Synthesis of tetrahydropyrazino[1,2-*a*]indole-1,4-diones

(Scheme 102) [168]. The starting indolenines (**494**) can be, in turn, easily obtained from phenylhydrazines and 2,2-dialkyl acetaldehydes following a Fischer protocol.

Following a one-pot two-step methodology, García-Valverde et al. described the diastereoselective synthesis of pyrrolpiperazine-2,6-diones (**502**). The synthesis is carried out through an Ugi/nucleophilic substitution/*N*-acylation/debenzoylation sequence using three doubly functionalised reactants –phenylglyoxal (**498**),  $\alpha$ -amino esters (**499**) and 3-bromopropionic acid (**468**)- along with the isocyanide (**31**; Scheme 103) [169].


Furthering the studies on the cyclisations of Ugi adducts promoted by the deprotonation of the amide nitrogen, Marcaccini's group proposed the synthesis of Ugi adducts with a highly electrophilic  $\alpha$ -acylamino substituent that would facilitate the intramolecular nucleophilic attack at this position. Therefore, they chose trichloroacetic acid (**448**) as one of the components of the U-4CC to facilitate the ring closure through an *N*-acylation leading to hydantoins (**505**). This synthesis is based on the enhanced electrophilic character of the trichloroacetamide group and the quality of trichloromethyl anion as leaving group, allowing the trichloroacetyl group to function as a masked carbonic acid surrogate. In

this way, the Ugi adducts (**503**) synthesised from primary amines (**114**), aryl aldehydes (**371**), trichloroacetic acid (**448**) and isocyanides (**31**), upon treatment with sodium ethoxide, underwent a rapid ring-closure reaction to give hydantoins (**505**) in good yields. Moreover, the products were easily isolated from the reaction medium by precipitation from the mother liquors (Scheme 104) [170].


This synthesis constitutes a more efficient methodology than the route described by Hulme's group for the synthesis of hydantoins (**509**) via an Ugi five-component condensation followed by a base-promoted cyclisation (Scheme 105) [171].

A similar strategy replacing trichloroacetic acid by propionic acid (**510**) has been reported by Chen et al. Surprisingly the base treatment of the Ugi adduct (**511**) affords the corresponding hydantoin (**509**), resulting from the behaviour of the acetylde anion as the leaving group. In contrast, the expected intramolecular Michael addition that would give rise to a diazepine did not occur (Scheme 106) [172].

The methodology described by Marcaccini's group has been employed for the synthesis of pseudopeptidic hydantoins (**516**) using Ugi/cyclisation/reduction/Ugi sequences, by inclusion of a nitro group as amine surrogate in the





Scheme 103 Synthesis of pyrrolpiperazine-2,6-diones




Scheme 104 Marcaccini's hydantoin synthesis

Scheme 105 Hulme's hydantoin synthesis






Scheme 106 Chen's synthesis of hydantoin



Scheme 107 Diastereoselective synthesis of pseudo-peptidic hydantoin



Scheme 108 Dehaen tandem synthesis of hydantoin-fused benzodiazepines


carbonyl component (**513**). Interestingly, the pseudo-peptidic hydantoin (**516**) were obtained in many cases with a high diastereoselectivity (Scheme 107) [173].





This strategy has also been used for the synthesis of fused heterocycles, following two different approaches. The first approach starts with the synthesis of the hydantoin scaffold with the proper functionalisation for the subsequent synthesis of the corresponding fused heterocycle. This methodology was chosen by Dehaen et al. for the synthesis of

hydantoin-fused triazolobenzodiazepines (**518**), using 2-azido-benzaldehyde (**517**), propargylamine (**324**), trichloroacetic acid (**448**) and isocyanides (**31**) as starting materials for the Ugi reaction. The treatment of the Ugi adducts with sodium ethoxide afforded the functionalised hydantoin intermediates, which, when subjected to intramolecular azide-alkyne cycloaddition (IAAC), afforded the expected hydantoin-fused benzodiazepine derivatives (**518**; Scheme 108) [166].

The second approach starts with the synthesis of cyclic imines followed by an Ugi-Joullié reaction, which introduces the trichloroacetic acid (**448**) for the base-induced cyclisation to the hydantoin-fused system. In this manner, Martens et al. described the synthesis of two different families of fused heterocycles: hydantoin fused with oxa(thia)zolidines (**524**) and 1,4-benzothiazines (**528**). The methodology developed for the synthesis of oxa(thia)zolidine-fused hydantoin (**524**) combines two multicomponent reactions, an Asinger and an Ugi-Joullié reaction (Scheme 109), whilst the synthesis of the second family (**528**) began with the condensation of 2-bromo-2-methylpropionaldehyde (**526**) and 2-aminothiophenol (**525**) affording

Scheme 109 Asinger/Ugi-Joullié reaction for the synthesis of oxa(thia)zolidine-fused hydantoin



**Scheme 110** Synthesis of benzothiazine-fused hydantoin**Scheme 111** Synthesis of diazepine-fused hydantoin**Scheme 112** Cyclisation of Passerini adducts affording oxazole-derived orthoamides**Scheme 113** Synthesis of 2,*N*-diarylglycines by the cleavage of Ugi adducts of glyoxylic acids

the 2*H*-1,4-benzothiazines (**527**) required for the ulterior Ugi-Joullié reaction (Scheme 110) [174].

Hydantoin fused with piperazines and diazepines have also been synthesised following this strategy. Thus, Nelson et al. used an *N*-Boc deprotection/Ugi-Joullié condensation/cyclisation sequence. In this sequence the trifluoroacetic acid (**530**) used as reagent in the deprotection of the amine group plays a double role, acting also as the acid component in the Ugi-Joullié reaction, and finally as a carbonic acid surrogate in the last stage of the synthesis (Scheme 111) [175].

Conversely, the intramolecular NH attack on Passerini three-component adducts (**537**) of trifluoromethylketones (**535**) gave rare orthoamides (**538**), the result being highly

dependent on the nature of the isocyanide, as well as the temperature and catalyst charge (Scheme 112) [176]. These pentacyclic orthoamides (**538**) are structurally related to those obtained by Marcaccini (**307**) by the cyclisation of P-3C-2C adducts of 6-oxo-4-thiocarboxylic acids through a similar mechanism (Scheme 56) [92].

A similar intermediate was proposed by Marcaccini et al. in the synthesis of 2,*N*-diarylglycines (**543**) from the basic treatment of Ugi adducts (**540**) synthesised from aromatic aldehydes (**371**), anilines (**114**),  $\alpha$ -ketoacids (**539**) and isocyanides (**31**; Scheme 113) [177].

These chemical results were not achieved when aliphatic aldehydes or amines were used or when aldehydes were

replaced by ketones. Based on these experimental results, the formation of the arylglycine derivatives was explained by the base-induced formation of imidazolidinone intermediates (**544**), which would undergo the deprotonation at C5 position. This key deprotonation step is not possible when ketones, aliphatic amines and/or aldehydes were employed in the Ugi condensation, since, either there is no hydrogen at the C5 position, or this is not acidic enough. Rearrangement of anion **545** would then lead to the intermediate ketene anions (**546**), which would give directly the  $\alpha$ -amino acid salts (**542**) via the addition of water or methanol, followed by the hydrolysis of the corresponding methyl esters (Scheme 114).


### Amine NH as nucleophile on adducts of the hydrazoic acid variant of the Ugi reaction

A variant of the classic Ugi reaction uses hydrazoic acid in place of the usual carboxylic acids to obtain  $\alpha$ -amino tetrazoles. Marcaccini took advantage of the nucleophilic


character of the  $\alpha$ -amino group on the adducts to design post-condensation transformations introducing the appropriate complementary electrophilic groups with one of the components of the U-4CC.

Thus, the Ugi condensation involving methyl *o*-formylbenzoates (**547**), amines (**114**), isocyanides (**31**), and hydrogen azide (**219**; prepared in situ from trimethylsilylazide and MeOH or from dimethylamine hydrochloride and sodium azide) gives a primary adduct (**548**) that undergoes a spontaneous 1,5-dipolar cyclisation to give the  $\alpha$ -amino tetrazole (**549**). Next, nucleophilic cyclisation affords isoindolinones (**550**). This latter reaction takes place spontaneously when benzyl or alkyl amines are used, and requires basic conditions with aromatic amines (Scheme 115) [178].

Hulme used a similar strategy, in which the ester moiety necessary for the post-condensation amidation is localised in the isocyanide (**551**). This reacts with carbonyl compounds (**153**), amines (**114**) and trimethylsilylazide (**175**) to give  $\alpha$ -amino tetrazoles (**552**), which are heated in methanol





**Scheme 114** Mechanism for the cleavage of Ugi adducts leading to 2,*N*-diarylglycines



**Scheme 115** Synthesis of tetrazoloisoindolines by post-condensation of azide-Ugi adducts

**Scheme 116** Synthesis of ketopiperazinotetrazoles by post-condensation of azide-Ugi adducts





**Scheme 117** Synthesis of tetrazolobenzodiazepines by post-condensation of azide-Ugi adducts

to give fused ketopiperazinetetrazoles (**553**, Scheme 116) [179].

Analogous tetrazolobenzodiazepines (**556**) were likewise synthesised by Voskressensky in a one-pot U-4CC of ketones (**153**), ammonium chloride (**132**), bifunctional isocyanides (**554**), and sodium azide (**555**) or trimethylsilylazide (**175**), followed by cyclisation (Scheme 117) [180]. Dömling used the same strategy, substituting primary amines (**114**) for ammonium chloride (**132**) [181]. In this case, ester hydrolysis followed by EDC/HOBt mediated amide bond formation was required to obtain the final tetrazolobenzodiazepines (**556**; Scheme 117).


Recently, Dömling prepared unsubstituted tetrazolo  $\gamma$ - and  $\delta$ -lactams (**560**) by the Ugi reaction of aliphatic ester-substituted aldehydes (**558**), trityl amine (**557**), isocyanides (**31**) and trimethylsilylazide (**175**), followed by deprotection with TFA and cyclisation with sodium hydride (Scheme 118) [182].

## Additional nucleophilic groups

### Nitrogen nucleophiles


Nucleophilic groups present in the components of the Ugi and Passerini reactions may interfere with the condensations, leading to unexpected reaction pathways (See, for example, the reaction between 2-hydroxybenzaldehyde (**700**), isocyanides (**31**) and ammonium formate (**121**;

Scheme 147) [183]. To avoid these competitive reactions, different strategies have been developed that use masked internal nucleophiles, such as *N*-protected amines or nitro or azide groups as amine surrogates. If these masked groups are properly located relative to complementary electrophilic sites, intramolecular reactions take place upon their activation, affording a variety of often pharmacologically relevant scaffolds. Thus, while Hulme et al. developed the UDC concept (Ugi reaction/Deprotection/Condensation) [184, 185], Marcaccini's group pioneered the use of nitro and azide groups as amine surrogates in Ugi/Condensation sequences. Thus, in a seminal work where the nitro group was introduced as a masked amino nucleophile in an IMCR, Marcaccini developed an elegant synthesis of furan derivatives (**563**) through Passerini/Knoevenagel sequences. When 2-nitrophenyl acetic acid (**561**) was chosen as the carboxylic component in a P-3CC with glyoxals (**333**) and isocyanides (**282**), the nitro-group played a double role, increasing the acidity of the benzylic position on the Passerini adduct to favour the intramolecular Knoevenagel condensation in mild conditions and acting as a masked amino-group. In this way, 2,5-dihydro-2-(2-nitrophenyl)-5-oxofuranes (**563**) were obtained upon treatment of the Passerini adducts (**562**) with piperidine and additional treatment with acid. Then, the reduction of the nitro-group afforded 2-oxoindoles (**565**) as a result of the nucleophilic attack of the amine on the lactone carbonyl group with subsequent opening of the lactone ring (Scheme 119) [186]. Thus, the overall reaction pathway involves a first intramolecular reaction with a carbon




**Scheme 118** Synthesis of tetrazolo lactams





Scheme 119 Marcaccini's three-step synthesis of 2-oxoindoles




Scheme 120 Marcaccini's synthesis of benzodiazepines by Ugi/reduction/cyclisation sequences

nucleophile, followed by a ring-switching transformation of a furan into an indole triggered by the attack of the initially masked nucleophile.

Later, Marcaccini's group described the synthesis of benzo[1,4]diazepine systems with different substitution patterns (**569**, **572**, **574**) through two-step Ugi/reduction/cyclisation (URC) sequences, varying the nature of the nitro derivative and/or the doubly functionalised reactant with the electrophilic site required for the cyclisation step.

Thus, the combination of 2-nitrobenzoic acid (**566**) with  $\alpha$ -amino esters (**567**) led to 1,4-benzodiazepine-2,5-diones (**569**; Scheme 120a) [187], while its combination with phenacylamine (**570**; Scheme 120b) [188] or arylglyoxals (**333**; Scheme 120c) [189] afforded stable 4,5-dihydro-3*H*-benzo[1,4]diazepin-5-ones (**572**, **574**). However, when arylglyoxals (**333**) were combined with 2-nitrobenzylamine (**583**), unstable 4-benzoyl-4,5-dihydro-3*H*-benzo[*e*][1,4]diazepines were obtained.



**Scheme 121** Synthesis of thienbenzodiazepines reported by Guo

Guo used a similar strategy to prepare thienbenzodiazepines (**582**) to test their antitumor activity as p53-MDM2 protein–protein interaction inhibitors. In this case, the use of convertible isocyanide (**576**) permits the obtention of benzodiazepine (**578**) by reduction of the nitro group on the non-isolated Ugi adduct (**577**). The treatment of **578** with Lawesson's reagent (**579**) affords compounds **580**, which can be further alkylated in the presence of DBU to give **582** (Scheme 121) [190].


Based on these results, García-Valverde et al. introduced new functionalised reagents to synthesise more complex systems. In this way, pyrrolbenzodiazepine and pyrroloquinazoline scaffolds have been synthesised from arylglyoxals (**333**) and amino or carboxylic reagents where the nitro group was attached [191, 192]. The reaction of glyoxals (**333**), 2-nitrobenzylamine (**583**), isocyanides (**31**) and 3-bromopropionic acid (**468**) and subsequent intramolecular nucleophilic attack of the peptidyl position onto the alkyl bromide leads to pyrrolidinones (**585**). Reduction with  $\text{SnCl}_2$  unmask the latent aromatic amine, resulting in pyrrolbenzodiazepines (**587**) or pyrroloquinazolines (**588**), depending on the reaction conditions (Scheme 122) [191].

Similarly, the same authors were able to synthesise pyrrolbenzodiazepines with different degrees of unsaturation (**590**, **591**, **593**) by the post-condensation transformation of arylglyoxals (**333**), isocyanides (**31**), 2-nitrobenzoic acid (**566**) and bifunctional amines (**589**, **324**, **592**; Scheme 123) [192].

Using a complementary approach, Marcaccini synthesised some analogous structures, as well as other medium size nitrogen heterocycles, from azide derivatives as amine surrogates, following Ugi/Staudinger/aza-Wittig cyclisation sequences. Thus, seven-membered rings as 4,5-dihydro-3*H*-benzo[1,4]diazepin-5-ones (**595**) were obtained from 2-azidobenzoic acid (**594**) and arylglyoxals (**333**) (Scheme 124, a) [193]; 5-oxo-4,5,6,7-tetrahydro-1*H*-1,4-diazepines (**597**) from 3-azidopropionic acids (**596**) and arylglyoxals (**333**; Scheme 124, b), and 2-oxo-1,4-benzodiazepines (**600**) from 3-phenyl-2-azidopropionic acid (**599**) and 2-aminobenzophenone (**598**; Scheme 124, c) [194]. Likewise, eight-membered rings as [(5*H*)-6-oxodibenzo[*b,f*][1,5]diazocines (**601**) were synthesised from 2-azidobenzoic acid (**594**) and 2-arylanilines (**598**; Scheme 124, d) [195].


The stereochemical outcome of these reactions have also been analysed, revealing uneven results. When the chiral information was incorporated into the acid component [194] the diastereoselectivity achieved was extremely poor, however high diastereoselectivities were observed when the chiral information was introduced in the amino component [187]. Moreover, reversal of diastereoselectivity was observed in the synthesis of 3-carboxamide-1,4-benzodiazepin-5-ones depending on the synthetic methodology employed [196].

Yan used aromatic amines linked to an azide group (**602**) to prepare benzimidazoles (**604**) or quinoxalin-2(1*H*)-ones (**606**) using an Ugi/Staudinger/aza-Wittig sequence. When




**Scheme 122** Three-step synthesis of pyrrolobenzodiazepines or pyrroloquinazolines developed by García-Valverde

**Scheme 123** Post-Ugi transformations to diverse pyrrolobenzodiazepines




carboxylic acids (**386**) were used in the Ugi reaction, the Ugi-adducts (**603**) reacted with triphenylphosphine at room temperature and successive heating in toluene to give benzimidazoles (**604**). In turn, when glyoxylic acids (**539**) were used, quinoxalin-2(1*H*)-ones (**606**) were formed at room temperature (Scheme 125) [197].

Wessjohann also employed the Ugi-Staudinger-aza-Wittig sequence to prepare imidazolines (**609**). Here, the Ugi adducts (**608**) of azidoalkylamines (**607**) were treated with triphenylphosphine or resin bound triphenylphosphine give the imidazolines (**609**). In some cases, *o*-amidines (**610**) were also observed (Scheme 126) [198].




**Scheme 124** Synthesis of 7- and 8-membered nitrogen heterocycles by Ugi/Staudinger/aza-Wittig cyclisation




**Scheme 125** Amino-azides in the Ugi- Staudinger-aza-Wittig process

Ding et al. have used analogous Ugi/Staudinger/aza-Wittig sequences using 2-azidobenzaldehyde (**611**) for the synthesis of different heterocycles. The reaction of 2-azidobenzaldehyde (**611**), amines (**114**), isocyanides (**31**), and carboxylic acids (**386**), followed by treatment

with methyl diphenyl phosphine yields the expected quinazolines (**613**; Scheme 127, a) [199]. When phenylglyoxylic acid (**376**) is used, 2-acylquinazolines (**615**) and/or 3*H*-1,4-benzodiazepin-3-ones (**616**) are obtained,




Scheme 126 Ugi-Staudinger-aza-Wittig synthesis of imidazolines reported by Wessjohann



Scheme 127 Ugi/Staudinger/aza-Wittig strategy employed by Ding

Scheme 128 Synthesis of indolo[1,2-c]quinazolines



depending on the steric and electronic nature of amine substituents (Scheme 127, a) [200].

When the Ugi-adducts (**617**) of 2-azidobenzaldehyde (**611**) with 2-acylaniline (**598**), isocyanides (**31**), and carboxylic acids (**386**) were treated with triphenylphosphine, the expected benzodiazocines (**621**) were not formed. Instead, indolo[1,2-*c*]quinazolines (**620**) were obtained, probably through quinazoline intermediates (**619**), which upon heating in toluene were transformed in situ by nucleophilic addition of the peptidyl carbon on the carbonyl group (Scheme 128). The authors argue that this is probably due to a restricted conformation of the iminophosphorane (**618**) that would be entropically unfavourable for the cyclisation between the iminophosphorane moiety and the ketone carbonyl group [201].


Ding has also used vinyliminophosphorane (**622**) as amine component in a U-4CC with 2-azidobenzaldehyde (**611**), isocyanides (**31**) and carboxylic acids (**386**). The resulting Ugi adducts (**623**) were treated with

triphenylphosphine and afforded the 3-arylidene-substituted 3*H*-1,4-benzodiazepines (**624**) by the usual Staudinger/aza-Wittig sequence (Scheme 129) [202].


Likewise, Ding carried out a tandem Passerini/Staudinger/aza-Wittig starting from  $\alpha$ -azidocinnamaldehydes (**625**), to obtain 2,4,5-trisubstituted oxazoles (**628**) in a one-pot process. A catalytic amount of potassium carbonate is necessary to the complete transformation of dihydrooxazoles (**627**) into the final oxazoles (**628**) *via* a 1,3-H shift (Scheme 130) [203]

Similarly, the hydrazoic variant of the Passerini condensation of 2-azidobenzaldehydes (**629**) in tandem with an acylation/Staudinger/catalytic aza-Wittig reaction readily gave 4-tetrazolyl-substituted 4*H*-3,1-benzoxazines (**631**; Scheme 131) [204].


Bazgir used 2-azidoacetic acid (**633**) in an Ugi reaction with 3-formylchromones (**632**), isocyanides (**31**) and amines (**114**). The Ugi adduct (**634**) was not isolated and was treated in situ with triphenylphosphine. When the reaction was carried out in water, the intermediate iminophosphorane was




Scheme 129 Synthesis of benzodiazepines developed by Ding



Scheme 130 Tandem Passerini/Staudinger/aza-Wittig synthesis of oxazoles



Scheme 131 Passerini/acylation/Staudinger/catalytic aza-Wittig synthesis of benzoxazines




**Scheme 132** Synthesis of coumarinpyrazinones and oxazines reported by Bazgir

hydrolysed and an intramolecular nucleophilic attack of the resulting amine, followed by a ring opening, took place to give the 3-oxo-1,4-diazepine-5-carboxamides (**635**). In turn, when dry toluene was used, the aza-Wittig reaction between the iminophosphorane and the amide group gave coumarinpyrazinones (**636**). The expected attack of the iminophosphorane on the chromone carbonyl group is probably not favoured due to the restricted conformation of intermediate (**634**) and it was not observed (Scheme 132) [205].


Marcaccini also used azines and semicarbazones as components for the Ugi condensation containing masked nucleophiles useful for post-condensation transformations.

The reaction between hydrazine hydrate (**637**) and oxo-compounds (**153**) affords the azine (**638**), which reacts with 2-benzoylbenzoic acid (**639**), and cyclohexyl isocyanide (**282**) to give the Ugi adduct (**640**). Acid hydrolysis of the hydrazone (**640**) liberates the hydrazine nucleophile and triggers an intramolecular cyclisation that results in 4-phenyl-1-(2*H*)phthalazinone-2-alkanoic acid amides (**642**; Scheme 133) [206].

Semicarbazones (**643**) were used for the first time in an Ugi reaction by Marcaccini for the synthesis of triazines (**648**). The reaction of semicarbazones (**643**) with isocyanides (**31**) and benzoyl-, or 4-methoxybenzoylformic acid



**Scheme 133** U-4CC with masked hydrazines



**Scheme 134** Use of semicarbazones in the Ugi reaction

(**644**) gave the expected Ugi adducts (**645**), which by treatment with sodium ethoxide resulted in a nucleophilic attack of the amide to the glyoxylic acid moiety leading to [1, 2, 4] triazines (**647**). Treatment of triazines **647** with diazomethane (**29**) led to the formation of the corresponding *O*-methyl derivatives (**648**; Scheme 134) [207]. Interestingly, compounds **647** and **648** constitute a new class of conformationally constrained pseudopeptides.


### Carbon nucleophiles

Carbon C–H centres contiguous to electron-withdrawing groups can act as internal nucleophiles in the presence of an appropriate base or, sometimes, spontaneously. This strategy has been extensively used by Marcaccini and other

researchers to facilitate post-condensation reactions of Ugi and Passerini adducts.

Thus, Bossio and Marcaccini described the preparation of tetrasubstituted furans (**653**) by a Passerini reaction of arylglyoxals (**333**), isocyanides (**31**), and cyanoacetic (**381**) [208] or arenesulfonylacetic acids (**649**) [209] followed by subsequent treatment with base and diazomethane (**29**). In this process, the Passerini adducts (**650**) were formed and isolated, and then are subjected to a base triggered intramolecular Knoevenagel condensation between the oxo group and the activated methylene to give the furan derivatives (**651**–**652**). Methylated derivatives (**653**) were obtained by treatment of compounds **652** with diazomethane (Scheme 135).

In order to extend this methodology to the preparation of pyrrole derivatives (**657**), these authors used arylglyoxal



**Scheme 135** Synthesis of tetrasubstituted furans





anils (**654**) in place of arylglyoxals (**333**). Arylgyoxal anils (**654**) were prepared from arylglyoxals (**333**) and anilines (**49**) and subjected to an Ugi condensation with cyanoacetic acid (**381**) and isocyanides (**31**). Cyclisation of the adducts (**655**) and methylation readily led to the desired pyrroles (**657**; Scheme 136) [124].

Analogously, the Ugi reaction of aldehydes (**81**), phenacylamine hydrochloride (**658**), cyanoacetic acid (**381**), and cyclohexyl isocyanide (**282**) affords the intermediate Ugi adducts (**659**), which cyclise spontaneously to give pyrrolinones (**660**). Subsequent treatment with diazomethane (**29**) yields methylated pyrroles (**661**; Scheme 137) [210].

More recently, Marcaccini, Torroba and Marcos performed U-4CC employing diarylgyoxal monohydrazones (**662**) as the amino components, isocyanides (**31**), aldehydes or ketones (**153**), and carboxylic acids bearing an activated methylene group, such as cyanoacetic acid (**381**), tosylacetic acid (**649**) and malonic acid monoethyl ester (**663**). The resulting Ugi products (**664a-c**) spontaneously cyclised via an intramolecular Knoevenagel condensation to 3(2*H*)-pyridazinones (**665a-c**; Scheme 138) [125].


The tandem Ugi–Knoevenagel strategy was also applied to the synthesis of quinolin-2-(1*H*)-ones (**668**), starting from 2-aminophenylketones (**666**), aldehydes (**81**),

**Scheme 136** Synthesis of tetrasubstituted pyrroles



**Scheme 137** Synthesis of *N*-cyclohexyl-2-aryl-2-(3-cyano-2-methoxy-4-phenylpyrrol-1-yl)acetamides

**Scheme 138** One-pot synthesis of pyridazinones




cyclohexylisocyanide (**282**) and malonic or arylsulfonyl-acetic acid derivatives (**381**, **649**, **663**; Scheme 139) [126].

In continuation with these post-Ugi transformations, Marcos et al. synthesised the Ugi adducts (**670**) of 3-formylchromones (**669**) with cyanoacetic acid (**381**), amines (**114**) and isocyanides (**31**). Treatment of these adducts (**670**) with KOH in methanol led to a ring-opening/ring-closing process that produced efficiently polyfunctionalised pyridones (**671**) related to cardiotoxic agent milrinone (Scheme 140) [127].


Going one step further, they developed a one-pot, two-step diastereoselective synthesis of spiropyrrolidinochromanones (**673**). The Ugi reaction between 3-formylchromones (**669**),

amines (**114**), isocyanides (**31**), and glyoxylic acids (**644**) affords the Ugi adducts (**672**) that suffer a nucleophilic conjugate addition to the position 2 of the chromone ring of a second amine (**114**) and subsequent intramolecular cyclisation to give the spiranic structures (**673**) in diastereoselective manner. Remarkably, three new stereogenic centers are formed in the cyclisation process (Scheme 141). Ugi adducts are stable and can be isolated, *but* their purification is not necessary and can be directly used in the following step [211].


On the other hand, Marcaccini et al. also reacted isocyanides (**31**) with cyanoacetic acid (**381**) in the absence



Scheme 139 Ugi/Knoevenagel strategy for the synthesis of quinolinones



Scheme 140 Synthesis of milrinone analogues



Scheme 141 Diastereoselective synthesis of spiropyrrolidinochromanones


of carbonyl compounds affording *N,N'*-disubstituted 2-cyanoacetoxy-2-cyanomethylmalondiamides (**681**). According to the mechanism proposed by the authors, the key step is the formation of intermediate ketene (**676**) by deamidation of the adduct of cyanoacetic acid and isocyanide (**674**). Cyanoketene (**676**) is then attacked by a second molecule of isocyanide (**31**) and cyanoacetic acid (**381**) to form the intermediate (**678**). The carbonyl group in (**678**) undergoes a regular Passerini reaction and eliminates a further molecule of cyanoketene (**676**), ensuring the cyclic continuation of the reaction. It is also possible that cyanoketene elimination takes place before the final Passerini condensation (Scheme 142) [212].

### Nucleophilic groups introduced with external reagents


The addition of, usually bifunctional, nucleophiles to Passerini and Ugi adducts containing complementary electrophilic sites can lead to the formation of novel heterocyclic structures in a very efficient manner. With this aim Marcaccini developed different post-condensation transformations using

ammonia or bisamine equivalents. For example, he treated arylglyoxals (**333**) with carboxylic acids (**386**) and isocyanides (**31**) to give *N*-alkyl-2-acyloxy-3-aryl-3-oxopropionamides (**682**). Upon heating of this Passerini adduct (**682**) with ammonium formate (**121**) in acetic acid, 5-oxazolecarboxamides (**684**) were obtained in fair yields (Scheme 143) [213, 214]. This strategy takes advantage of the reactivity of the peptidyl carbon on intermediate (**683**) and is in that way similar and complementary to the synthesis of oxazoles from Ugi adducts shown in Scheme 90. Interestingly, when arylthioacetic acids (**386**,  $R^3 = \text{CH}_2\text{SAr}$ ) were used, sulfur-substituted oxazole-5-carboxamides (**684**) were obtained [215].


Taking this idea further, the Ugi reaction between arylglyoxals (**333**), amines (**114**), benzoylformic acid (**376**) and isocyanides (**31**) afforded the Ugi adducts (**685**). The four-component reaction was carried out in two different manners: preforming the corresponding imines (**654**) from amines (**114**) and arylglyoxals (**333**) or following the classical one-pot procedure. The two methods led to the expected product with similar yields as an equilibrium mixture of tautomers (**685a** and **685b**). Treatment of these Ugi adducts (**685**) with



Scheme 142 Reaction of isocyanides with 2-cyanoacetic acid



Scheme 143 Synthesis of oxazole-5-carboxamides developed by Marcaccini




**Scheme 144** Synthesis of phenylpyrazine-2-carboxamides

ammonium acetate (**188**) afforded 1,*N*-disubstituted 4-aryl-1,6-dihydro-6-oxo-5-phenylpyrazine-2-carboxamides (**686**) by a Davidson type cyclisation (Scheme 144) [216].

Marcaccini also described the synthesis of benzothiazepinones (**691**) from 2-chloro-5-nitrobenzaldehyde (**687**), amines (**114**), isocyanides (**31**), and chloroacetic acid (**359**). Treatment of the Ugi adduct (**688**) with thiourea (**689**) resulted in the nucleophilic substitution of the aliphatic chlorine atom, followed by an aromatic nucleophilic substitution to give the benzothiazepinones (**691**). Then, the nitro-group could be reduced with iron in acetic acid to give an amine (**692**) that reacted with phenyl isothiocyanate (**693**) resulting in thiourea (**694**) (Scheme 145) [217].


### Adduct cleavage

Cleavage reactions of the IMCR adducts can lead to compounds difficult to obtain by other procedures. An example of this strategy used by Marcaccini for the synthesis of glycine derivatives (**543**) is shown in Scheme 113. Using a different approach, Marcaccini and Marcos synthesised different 1,3-dicarbonyl compounds in two steps by a Passerini/hydrolysis or a Passerini/reduction strategy. The reaction between glyoxals (**695**), isocyanides (**31**) and acetic acid (**224**) gave the expected Passerini adducts (**696**). When these  $\alpha$ -acyloxyamides (**696**) were treated with activated zinc in aqueous methanol,



**Scheme 145** Synthesis of 4,5-dihydro-1,4-benzothiazepin-3(2*H*)-ones

**Scheme 146** Post-Passerini transformations leading to 1,3-dicarbonylic compounds




the corresponding  $\beta$ -keto amides (**697**) were obtained (Scheme 146, a) [218]. When glyoxylamides or glyoxyl esters (**695**,  $R^1 = \text{NHR}$  or  $\text{OR}$ ) were used in the Passerini condensation, tartronodiamides (**698**,  $R^1 = \text{NHR}$ ) and tartronoamidoesters (**698**,  $R^1 = \text{OR}$ ) were obtained by zinc catalysed solvolysis of the corresponding Passerini adducts (**696**; Scheme 146, b), [219]. Finally, when the Passerini adducts of glyoxylamides (**696**,  $R^1 = \text{NHR}$  or  $\text{NRR}'$ ) were reduced with samarium diiodide, malonodiamides (**699**) were obtained (Scheme 146, c) [220]. This protocol permits, in a simple and efficient manner, the synthesis of malonic subunits that can be useful in the synthesis of retro-peptides.

### Ugi reactions with phenols



Marcaccini pioneered the use of phenols as substitutes for carboxylic acid in Ugi reactions [183]. Reaction of salicylaldehyde (**700**) with ammonium formate (**121**) and isocyanides (**31**) readily gave benzofuran derivatives (**703**; Scheme 147). In this case, the primary adduct (**702a**) is obtained through an intramolecular attack of the phenoxide on the nitrile cation in the reaction intermediate (**701**). The tautomerisation of this primary adduct produces diamino-furan (**702b**), which reacts with a second molecule of salicylaldehyde (**700**) giving benzofuran (**703**).

Several years later, Kobayashi reported the preparation of 2,3-bis(arylamino)benzofurans (**706**) and/or 2,3-diimino-2,3-dihydrobenzofurans (**707**) based on the reaction




**Scheme 147** Synthesis of benzofurans by an intramolecular phenol-Ugi reaction

**Scheme 148** Kobayashi's synthesis of diaminobenzofurans based on Marcaccini's chemistry




**Scheme 149** González-Zamora approach for the synthesis of 2-imino-1,4-benzoxazines



**Scheme 150** Wada's synthesis of benzoxazoles

**Scheme 151** Ukaji's synthesis of iminobenzothiophenones



of 2-aryliminophenols (**704**) and aryl isocyanides (**705**) with boron trifluoride catalysis (Scheme 148) [221]. A similar approach, using a secondary amine, electron-poor salicylaldehydes and aliphatic or aromatic isocyanides in the presence of catalytic  $\text{SiO}_2$  was reported by Ramazani [222]. In this latter case, the presence of a tertiary amine on benzofuran position 3 does not allow further oxidation to the corresponding diimine derivative and the 2,3-diaminobenzofurans are isolated as the only products. Very similar synthetic strategies were later published by Chattopadhyay's [223] and Adib's [224] groups, using respectively cerium ammonium nitrate as Lewis acid catalyst and water as solvent.

A related application, in which two phenol groups compete as acid components in an intramolecular reaction, was

developed by González-Zamora et al. for the synthesis of 2-imino-1,4-benzoxazines (**712**; Scheme 149) [225].

Wada et al. published a synthesis of 2-(alkoxyalkyl)benzoxazoles (**717**) by a tandem migration/carboalkoxylation of *o*-isocyanophenyl acetals (**713**) catalysed by  $\text{BF}_3 \cdot \text{OEt}_2$  and 2,4,6-collidine. The authors propose two alternative mechanisms for this transformation, one of which resembles a Passerini-type reaction with an ethylideneoxonium intermediate (**716**) and the phenol group acting as acid component (Scheme 150) [226].

The reaction of thiophenols (**718**) as acid components in intramolecular Passerini-type reactions has also been recently reported by Ukaji et al. (Scheme 151) [227]. The reaction is promoted by  $\text{LiI}$ , and in situ oxidation of the


Passerini adduct (**719**) leads to benzothiophene-3(2*H*)-one (**720**).

Shiri et al. performed a similar reaction starting from 2-mercaptoquinoline-3-carbaldehyde (**721**), resulting in tricyclic 2-(cyclohexylimino)thieno[2,3-*b*]quinolin-3(2*H*)-ones (**722**) through an intramolecular Passerini/oxidation tandem process (Scheme 152) [228].


An interesting related precedent had been published by Marcaccini 20 years earlier. Thus, Marcaccini et al. allowed 2-(arylaminothiocarbonyl)cyclohexanones (**723**) and isocyanides (**31**) to react in an acidic medium to

give novel diimino thioanhydrides (**728**; Scheme 153). Although the reaction can be considered as a formal [4 + 1] cycloaddition, the authors propose a two-step mechanism involving an initial protonation of carbonyl group, followed by the nucleophilic attack of the carbenoid carbon of the isocyanide and nucleophilic attack of the thioamide sulfur on the intermediate imidoyl cation (**726**). Final elimination of water takes place easily because of the highly conjugated nature of the final products (**728**). Salicylic acid (**724**) was found to provide a suitable acid catalysis to facilitate this process [229].


**Scheme 152** Shiri's synthesis of iminothiophenone-fused quinolines



**Scheme 153** Formal [4 + 1] cycloaddition 3-oxothioamides and isocyanides



**Scheme 154** Ugi-Smiles reaction developed by El Kaïm




Later El Kaim developed a general-purpose strategy in which a condensation of Ugi with phenols (**729**) is followed by a Smiles rearrangement, resulting in four-component adducts (**734**) in good yields (Scheme 154) [230, 231]. The reaction takes place with electron-deficient phenols, typically ortho or para-nitrophenols (**729**). The primary adduct (**732**) undergoes a Smiles rearrangement to give rise to nitrophenyl aminoamides (**734**).

This strategy certainly increases the diversity of structures obtainable in Ugi-type reactions, although it is limited to the use of a relatively small number of phenols containing electron-withdrawing substituents.


## Enol-Ugi reactions

Enols have  $pK_a$  values comparable to that of phenols, while enolates are more nucleophilic than carboxylates. In 2012 Marcaccini and Marcos showed that heterocyclic enols (**736**) are feasible acid components in Ugi-type condensations [232]. The presence of an  $\alpha,\beta$ -unsaturated electron-withdrawing group in the enol is a structural determinant that facilitates a Michael-retro-Michael rearrangement of the primary adduct (**739**; Scheme 155). This constitutes the driving force of the reaction and


**Scheme 155** Mechanism of the enol-Ugi reaction




**Fig. 3** Heterocyclic enamines obtained by the enol-Ugi reaction







Scheme 156 Charton's synthesis of symmetrical squaramides



Scheme 157 Ramazani's enol-Ugi synthesis of tropolone-derived enamines



affords the heterocyclic enamine products (741). Different 5- and 6-membered enolic heterocycles can be used, including pyrrolidine-2,3-diones, furane-2,3-diones, 3- and 4-hydroxycoumarins and dihydropyrimidines, resulting in different heterocyclic enamines (742–746; Fig. 3) [232–234].

Independently and almost simultaneously, Julie Charton's group published a related double 4-component Ugi-type reaction using squaric acid (747) as the acid component (Scheme 156) [235]. This reaction is also possible when substituting squaramic or squaramide acids for squaric acid, as it was later demonstrated by Mehrabi [236].

On the other hand, Ramazani et al. successfully used tropolone (749) as the enol component in enol-Ugi reactions leading to 2-(*N,N*-dialkylamino)-2,4,6-cycloheptatrien-1-one derivatives (750; Scheme 157) [237].

More recently, Rabêlo and Echemendía described an enol-Ugi reaction between 2-hydroxy-3-nitro-1,4-

Scheme 158 Rabêlo and Echemendía's synthesis of substituted naphthoquinones



Scheme 159 Ugi-type condensation with saccharin

naphthoquinone (752), different secondary diamines (751) and isocyanides (31) to give 3-substituted 1,4 naphthoquinones (753; Scheme 158) [238].


In addition, Ramezanpour et al. reported a related variant of the Ugi condensation using saccharin (755) as acid surrogate (Scheme 159) [239].

In line with previous research, Marcos et al. developed analogous enol-Passerini and pseudo-enol-Ugi reactions from pyrrolidinodiones (757), isocyanides (31) and aldehydes (81; Scheme 160) [240].

On the other hand, the post-condensation transformation of hydroxycoumarin enol-Ugi adducts (762) led to an efficient synthesis of geometrically restricted peptidomimetic chromeno[3,4-*b*]pyrazin-5-ones (763; Scheme 161) [241].

Paixão et al. have developed an elegant stereoselective synthesis of tetrahydropyridines (769) [242] and related natural product hybrids (774) based on an intramolecular enol-Ugi condensation (Scheme 162) [243]. Interestingly, the reaction of hemiacetal (767 or 772) and an amine (114 or 773) generates a reactive intermediate (768) containing the enol and Schiff base moieties that make possible the ensuing enol-Ugi condensation.

**Scheme 160** Enol-Passerini and pseudo-enol-Passerini reactions



**Scheme 161** Chromenopyrazinone synthesis by an enol-Ugi post-condensation transformation

## Cycloaddition reactions of isocyanides


### [3 + 2] Cycloadditions of bifunctional isocyanides

Isocyanoacetate derivatives (**1**) and other bifunctional isocyanides have been used by Marcaccini for the synthesis of heterocycles by  $\alpha$ -addition to the isocyanide carbon and subsequent cyclisation on the acidic  $\alpha$  position, as described earlier in this review. These bifunctional isocyanides can also be used in formal [3 + 2] cycloaddition reactions with dipolarophiles to afford 5-member nitrogen heterocycles.


The variation of the van Leusen reaction employing activated alkenes, such as  $\alpha,\beta$ -unsaturated ketones, esters or nitriles, was reported in 1972 for pyrrole synthesis [244]. However, it was not until early 1990s, when Barton et al. made aryl nitroolefins react with TOSMIC in

tetrahydrofuran-isopropanol, at  $-78\text{ }^\circ\text{C}$ , in the presence of DBU, to afford 3-nitropyrroles (Scheme 163) [245]. The formal [3 + 2] cycloaddition probably involves an asynchronous mechanism consisting in a first Michael addition of TOSMIC  $\alpha$  carbon to the nitroolefin, followed by internal attack of the nitronate on the isocyanide group. Proton exchange, elimination of the sulphone and aromatisation through a [1, 5] sigmatropic shift of hydrogen leads to the formation of a 3-nitropyrrole (**782**) in rather poor yields. The yields were though significantly improved by Ono et al., who carried out the reaction with NaH in DMSO [246].

Marcaccini's group applied this knowledge to the synthesis of sugar derivatives from TOSMIC sodium salt (**784**) and nitro-glucoside derivatives (**783**). In this way, nitropyrroles substituted with sugar functionalities were successfully obtained (Scheme 164) [247].



Scheme 162 Intramolecular enol-Ugi condensation by Paixão and Rivera






Scheme 163 Barton's synthesis of 3-nitropyrroles

In a study contemporary to Marcaccini's research, Van Leusen et al. also reported the synthesis 3-nitropyrroles from TOSMIC and nitrostyrenes, in almost quantitative yield and high-regioselectivities (Scheme 165) [248].

Later research by Krishna et al. expanded Marcaccini studies by reacting TOSMIC and activated alkene bonds to

different ribose derivatives to obtain 3-nitropyrrole C-nucleosides (Scheme 166) [249, 250].

A different approach was developed by Marcaccini, based on the generation of a novel class of nitrile ylides by the treatment of isothiocarbamoyl chlorides (792) with triethylamine. These nitrile ylides react readily with dienophiles as

**Scheme 164** Synthesis of C-glycosyl pyrroles**Scheme 165** Van Leusen synthesis of 3-nitropyrroles**Scheme 166** Synthesis of pyrrole C-Nucleosides

acetylenedicarboxylate (**794**) through a 1,3-dipolar cycloaddition to give 2-*H*-pyrroles (**795**). The precursor isothiocarbamoyl chloride (**792**) was, in turn, easily prepared by the reaction of sulfenyl chlorides (**790**) and 2-isocyanopropionitrile (**791**; Scheme 167) [251].


This strategy was extended to the cycloaddition of nitrile ylides (**798**) obtained from 4-nitrobenzoylisocyanides (**796**) and arenylsulfenyl chlorides (**790**). In this way, different


1-*H*-pyrroles (**800**) and 1-*H*-imidazoles (**801**) were readily prepared by the reaction with, respectively, dimethyl acetylenedicarboxylate (**794**) or ethyl cyanofornate (**799**; Scheme 168) [252].

### [4 + 1] Cycloadditions of isocyanides


The  $\alpha$ -addition of 1,4-bifunctional reagents to isocyanides can result in 5-member heterocycles, product of a formal [4 + 1] cycloaddition process. Saegusa found that stoichiometric Et<sub>2</sub>AlCl promotes the [4 + 1] cycloaddition of isocyanides with  $\alpha,\beta$ -unsaturated carbonylic compounds to afford unsaturated N-substituted iminolactones [253]. Later, this reaction was effectively carried out by Chatani in the presence of catalytic GaCl<sub>3</sub> [254]. Yavari has similarly synthesised 2-aminofurans [255], although, several reports point out the instability of 2-aminofurans and their tendency to be quickly oxidized in the reaction medium [256]. On the other hand, Marcaccini and Marcos implemented different strategies to trap the elusive 2-aminofurans obtained by the [4 + 1] cycloaddition of isocyanides, resulting in tandem processes leading to diverse heterocyclic systems.

Thus, an yttrium triflate catalysed tandem process involving the [4 + 1] cycloaddition of isocyanides (**31**) with  $\alpha,\beta$ -unsaturated ketoester (**802**) and the ensuing [4 + 2] cycloaddition of the resulting unstable aminofuran (**804**) with maleimide derivatives (**803**) successfully gave polysubstituted anilines (**806**) in moderate to good yields (Scheme 169) [257]. The intermediate oxabicyclic (**805**) was not isolated, but opening of the oxygen bridge and dehydration take place immediately in the reaction medium.


**Scheme 167** 1,3-Dipolar cycloaddition of nitrile ylides derived from 2-isocyanopropionitrile



**Scheme 168** Synthesis of 1*H*-pyrroles and 1*H*-imidazoles via 1,3-dipolar cycloaddition



**Scheme 169** Synthesis of polysubstituted anilines by tandem [4+1]/[4+2] cycloaddition of isocyanides




**Scheme 170** [4+1]/[4+2] Cycloaddition strategy for the synthesis of 4-aminoxanthenes


Marcos et al. applied this methodology to a highly efficient synthesis of 4-aminoxanthenes (**808**), starting from 3-carbonylchromones (**807**), isocyanides (**31**) and

dienophiles (**803**; Scheme 170) [258]. The geometrical rigidity of the  $\alpha,\beta$ -unsaturated carbonyl of the chromone drives the progress of the reaction, which proceeds without the need for catalysis at near room temperature.

Interestingly, the use of non-symmetrical dienophiles, such as methylvinylketone or acrylonitrile (**809**), allows the isolation of the intermediate 1-hydroxydihydroxanthenes (**810**), which in this case turn out to be stable products that can be further dehydrated in the presence of a base to give the corresponding xanthenes (**808**; Scheme 171) [259, 260]. The reaction is fully regioselective so that the electron-withdrawing group  $R^2$  occupies the 3-position of the product xanthone or dihydroxanthone.



**Scheme 171** Synthesis of dixydroxanthones and xanthones



**Fig. 4** Secalonic acid analogue obtained by a tandem [4+1]/[4+2] cycloaddition of isocyanides

Xanthones related to natural products, such as secalonic acid analogue **809**, could be obtained using this simple procedure (Fig. 4).

A virtually identical synthesis of 4-aminoxanthones was later reported by Teimouri [261], who also used quinones as dienophiles to obtain 4 or 5 fused polycycles [262].

Benzocoumarins (**811**) were also obtained by Marcos et al. by the tandem [4+1]/[4+2] cycloaddition of


isocyanides (**31**), 3-carbonylcoumarins (**810**) and dienophiles (**803**; Scheme 172) [263].


Tang et al. have used *o*-alkenyl arylisocyanides (**813**), which react with  $\alpha,\beta$ -unsaturated ketones (**812**) in a tandem [4+1]/intramolecular [4+2] process to afford carbazole and indolocarbazole alkaloid derivatives (**814**; Scheme 173) [264]. This strategy was recently used for the synthesis of the alkaloid malasseziazole C **815** [265].

A different approach was also developed by Marcos' group, where polycyclic isoindoles (**818** and **819**) were obtained in a tandem multicomponent synthesis starting from cyclic 1,3-dicarbonyls (**816** or **817**), aldehydes (**81**), isocyanides (**31**), and maleimides (**803**). The reaction involves a sequence of a Knoevenagel condensation, and [4+1] and Diels–Alder cycloadditions. Interestingly, a further microwave-promoted dehydrogenative N–C bond forming reaction allows the synthesis of a natural product-like isoindolocarbazoles (**820**, Scheme 174) [266].


A related tandem Knoevenagel/double cycloaddition reaction was developed by Byk to obtain staurosporine analogues (**824**) [267]. This multicomponent reaction begins with the condensation of aldehyde (**81**) and substituted chiral tetramic acid (**821**), which under the reaction conditions suffer successive cycloadditions with an

**Scheme 172** Tandem synthesis of benzocoumarins






Scheme 173 Tandem [4+1]/intramolecular [4+2] cycloaddition for the synthesis of carbazoles



Scheme 174 Tandem synthesis of polycyclic isoindoles



Scheme 175 Byk's synthesis of isoindoles


isocyanide (**31**) and a dienophile (**823**) to give the desired isoindoles (Scheme 175).

An intramolecular version of this type of process was again reported by the group of Tang in 2020, in which polyfunctionalised cyclo[*b*]fused carbazoles (**826**) are obtained in a catalyst-free aqueous media with yields up to 99% (Scheme 176) [268]. Di- and tricarbazoles (**827**) were obtained from di- and trialdehydes.


A related strategy, developed by Xu's group consists in the formation of an intermediate furane (**831**) by the addition of alkenyl isocyanides (**828**) to ene-yne-ketones (**829**)

and a subsequent intramolecular Diels–Alder cycloaddition (Scheme 177) [269].


In the absence of a dienophile, the [4+1] cycloaddition of formylchromones (**833**) and isocyanides (**31**) gives unstable aminofuranes (**834**) that can undergo a Friedel–Crafts-type addition to a second molecule of the aldehyde (**833**) and dehydrate to give chromenylmethylene furochromones (**835**; Scheme 178). Preliminary studies of this reaction were reported by the group of Marcos in 2008 [270, 271] and later published a complete study by spectroscopic and X-ray diffraction analyses in which the structure (**835**) was



**Scheme 176** Tandem Knoevenagel/[4+1]/intramolecular [4+2] cycloaddition to cyclo[*b*]fused carbazoles



**Scheme 177** Tandem 1,6-addition/cyclisation/[4+2] cycloaddition



**Scheme 178** Tandem [4+1] cycloaddition/condensation for the synthesis of chromenylmethylene furochromones

unequivocally confirmed [272]. Meanwhile, a report by Bandyopadhyay in 2010 [273] published the same reaction, but postulated a structure for the products that was later shown to be incorrect [274]. Teimouri also reported the same reaction with similar results [275].

The Friedel–Crafts trapping of 3-formylchromone derived aminofuranes with different electrophiles, such as azodicarboxylates (**836**) [276], isatines (**838**) [277] and arylidene malononitriles (**840**) [278], was later reported by Teimouri et al. (Scheme 179).


They also performed the reaction of 3-formylchromones (**833**), isocyanides (**31**) and carboxylic acid anhydrides (**842**), which led to the formation of (acyloxymethylidene)

chromonyl-furochromones (**844**) by the Friedel–Crafts attack of intermediate aminofurane (**834**) to an *O*-acylated chromone (**843**; Scheme 180) [279].


On the other hand, chromone 3-carboxylic acid (**845**), undergoes [4+1] cycloaddition with isocyanides (**31**) to give a highly reactive intermediate iminoanhydride (**846**), which in the presence of alcohol nucleophiles is opened to give chromone-2-carboxamido-3-esters (**847**; Scheme 181). If water is used as nucleophile, the resulting carboxylic acid decarboxylates spontaneously to give 2-amido-substituted chromanones (**848**) [280].

The product chromone-2-carboxamido-3-esters (**847**) are obtained as a mixture of the *cis* (**847cis**) and enol (**847en**)





Scheme 179 Tandem [4+1]/Friedel-Crafts reactions




Scheme 180 1,4-addition of intermediate aminofuranes to acylated formylchromones

tautomers, which equilibrate in solution to a thermodynamic mixture of *cis* (**847cis**), *trans* (**847trans**) and enol (**847en**) isomers (Scheme 182).



A related-isocyanide four-component reaction was published by Shaabani in 2008 using Meldrum's acid to obtain coumarin 4-carbamoyl-2-oxochromane-3-carboxylates (**852**) [281]. The mechanism involves the [4+1]

cycloaddition of isocyanide (**31**) with the  $\alpha,\beta$ -carbonyl intermediate, formed in situ through the condensation of aldehyde (**849**) and the Meldrum's acid (**850**; Scheme 183). In this case, the major isomer was **852cis**, especially when bulky alcohols (**851**) and isocyanides (**31**) were used.



**Scheme 181** Conjugate addition of isocyanides to chromone 3-carboxylic acid

**Scheme 182** Equilibrium of isomers in chromone-2-carboxamido-3-esters



**Scheme 183** Four-component synthesis of 4-carbamoyl-2-oxochromane-3-carboxylates

transformations of adducts of classic multicomponent reactions, such as P3CC and U4CC, and development of new isocyanide multicomponent reactions using intramolecular methodologies and alternative reagents like phenols and enols.

With this review we want to gather the most important contributions of Stefano Marcaccini to organic synthesis and pay a fitting tribute to his memory.

**Acknowledgements** We wish to thank Prof Roberto Pepino, Dr Cristina Faggi, Dr Gloria Menchi and Beatrice Marcaccini for their help with this review.

**Author contributions** The authors declare that they have no competing financial interests or personal relationships that could have influenced this paper. All authors contributed equally to this work.

**Funding** Open Access funding provided thanks to the CRUE-CSIC agreement with Springer Nature.

## Declarations

**Competing interest** The authors declare that they have no competing financial interests or personal relationships that could have influenced this paper.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are

## Conclusions

Stefano Marcaccini is recognised as one of the foremost pioneers in isocyanide chemistry. The profound impact of his developments is clearly visible today in the work of those who follow his path and in the contributions of many first-rate research groups working in organic synthesis.

This review collects some of the overwhelming body of work developed by Stefano Marcaccini in the field of isocyanide chemistry. His encyclopaedic knowledge and his great creativity, as well as his scientific acumen, allowed him to utilise different strategies for the synthesis of heterocycles, some of which were still unpublished structures. Among other advancements, he made important contributions in the use of difunctional isocyanides, cycloaddition reactions of isocyanides, post-condensation

included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

- Schröder R, Schöllkopf U, Blume E, Hoppe I (1975) Synthesen mit  $\alpha$ -metallierten isocyaniden, xxviii. In 2-stellung unsubstituierte oxazole aus  $\alpha$ -metallierten isocyaniden und acylierungsreaktionen. *Justus Liebigs Ann Chem* 1975:533–546. <https://doi.org/10.1002/jlac.197519750315>
- Schöllkopf U (1977) Recent applications of alpha-metalated isocyanides in organic synthesis. *Angew Chem Int Ed* 16:339–422. <https://doi.org/10.1002/anie.197703393>
- van Leusen D, van Leusen AM (2001) Synthetic use of tosylmethyl isocyanide (TosMIC). In: Overman LE (ed) *Organic reactions*. Wiley, New York, pp 417–666
- Bossio R, Marcaccini S, Pepino R (1986) A novel synthetic route to oxazoles: One pot synthesis of 2-arylthio-5-alkoxyoxazoles. *Heterocycles* 24:2003–2005. <https://doi.org/10.3987/R-1986-07-2003>
- Bossio R, Marcaccini S, Pepino R, Polo C, Torroba T (1991) A novel class of oxazole derivatives-4-acyl-2-arylthio-5-ethoxyoxazoles. *Org Prep Proced Int* 23:670–672. <https://doi.org/10.1080/00304949109457925>
- Gulevich AV, Zhdanko AG, Orru RVA, Nenajdenko VG (2010) Isocynoacetate derivatives: synthesis, reactivity, and application. *Chem Rev* 110:5235–5331. <https://doi.org/10.1021/cr900411f>
- Bossio R, Marcaccini S, Pepino R (1986) Synthesis of 5,5'-dialkoxy-2,2'-dioxazolylsulfides. *Heterocycles* 24:2411–2413. <https://doi.org/10.3987/R-1986-09-2411>
- Bossio R, Marcaccini S, Pepino R, Torroba T, Valle G (1987) An unusual and simple one-pot synthesis of thiazolo[5,4-*d*]thiazoles. *Synthesis*. <https://doi.org/10.1055/s-1987-28202>
- Rössler A, Boldt P (1998) Improved access to thiazolo[5,4-*d*]thiazole and thieno[2,3-*d*]thiazole. *J Chem Soc Perkin Trans 1*(1998):685–687
- Reginato G, Mordini A, Zani L, Clamante M, Dessi A (2016) Photoactive compounds based on the thiazolo[5,4-*d*]thiazole core and their application in organic and hybrid photovoltaics. *Eur J Org Chem* 2016:233–251. <https://doi.org/10.1002/ejoc.201501237>
- Bossio R, Marcaccini S, Muratori M, Pepino R, Valle G (1990) Studies on alkyl isocynoacetates and related compounds - synthesis of 6-arylthio-8-ethoxycarbonyl-4-ethoxycarbonylmethylaminoimidazo[5,1-*b*][1,3,5]thiadiazine-2-thiones. *Heterocycles* 31:611–615. <https://doi.org/10.3987/COM-89-5243>
- Mampuy P, Zhu Y, Sergeyev S, Ruijter E, Orru RVA, Van Doorslaer S, Maes BUW (2016) Iodide-catalyzed synthesis of secondary thiocarbamates from isocyanides and thiosulfonates. *Org Lett* 18:2808–2811. <https://doi.org/10.1021/acs.orglett.6b01023>
- Guan Z, Zhu S, Wang S, Wang H, Wang S, Zhong X, Bu F, Cong H, Lei A (2021) Electrochemical oxidative carbon-atom difunctionalization: towards multisubstituted imino sulfide ethers. *Angew Chem Int Ed* 60:1573–1577. <https://doi.org/10.1002/anie.202011329>
- Xiaofang Lei YW, Fan E, Sun Z (2019) In situ activation of disulfides for multicomponent reactions with isocyanides and a broad range of nucleophiles. *Org Lett* 21:1484–1487. <https://doi.org/10.1021/acs.orglett.9b00275>
- Neo AG, Carrillo RM, Marcos CF (2011) A straightforward synthesis of 2-aminobenzothiazoles from Herz compounds. *Org Biomol Chem* 9:4850–4855. <https://doi.org/10.1039/C1OB05398A>
- Ugi I, Werner B, Dömling A (2003) The chemistry of isocyanides, their multicomponent reactions and their libraries. *Molecules* 8:53–66. <https://doi.org/10.3390/80100053>
- Bossio R, Marcaccini S, Pepino R (1995) Studies on isocyanides - synthesis of *N*-tosylguanidines. *Tetrahedron Lett* 36:2325–2326. [https://doi.org/10.1016/0040-4039\(95\)00246-9](https://doi.org/10.1016/0040-4039(95)00246-9)
- Bossio R, Marcaccini S, Pepino R (1995) Studies on isocyanides and related compounds - synthesis of 3- substituted 2-arylsulfonylamino-3,4-dihydro-4-oxoquinazolines via a novel 3-component reaction. *J Heterocycl Chem* 32:1115–1116
- Giustiniano M, Basso A, Mercalli V, Massarotti A, Novellino E, Tron GC, Zhu J (2017) To each his own: isonitriles for all flavors. Functionalized isocyanides as valuable tools in organic synthesis. *Chem Soc Rev* 46:1295–1357. <https://doi.org/10.1039/c6cs00444j>
- Bossio R, Marcaccini S, Pepino R, Polo C, Valle G (1989) A novel synthetic route to imidazole derivatives - synthesis of mesoionic 3-alkyl-2-arylthio-1,3-diazolium-4-olates. *Synthesis* 1989:641–643. <https://doi.org/10.1055/s-1989-27347>
- Garcia-Valverde M, Marcaccini S, Gonzalez-Ortega A, Rodriguez FJ, Rojo J, Torroba T (2013) Complementary regioselectivity in the synthesis of iminohydantoins: remarkable effect of amide substitution on the cyclization. *Org Biomol Chem* 11:721–725. <https://doi.org/10.1039/c2ob27098f>
- Bossio R, Marcaccini S, Pepino R, Polo C, Torroba T, Valle G (1989) Synthesis of 1-arylthiocarbonyl-4-isopropylamino-2,5-dihydro-1*H*-imidazole-2-thiones, a novel class of imidazole derivatives. *Heterocycles* 29:1843–1847. <https://doi.org/10.3987/COM-89-5089>
- Bossio R, Marcaccini S, Pepino R, Polo C, Torroba T (1989) Synthesis of 2,4-diarylthio-5-*N*-alkyl-*N*-phenylaminooxazoles: a novel class of oxazole derivatives. *Heterocycles* 29:1829–1833. <https://doi.org/10.3987/COM-89-5070>
- Sun X, Janvier P, Zhao G, Bienaymé H, Zhu J (2001) A novel multicomponent synthesis of polysubstituted 5-aminooxazole and its new scaffold-generating reaction to pyrrolo[3,4-*b*]pyridine. *Org Lett* 3:877–880. <https://doi.org/10.1021/ol007055q>
- Janvier P, Bois-Choussy M (2003) A one-pot four-component (ABC<sub>2</sub>) synthesis of macrocycles. *Angew Chem Int Ed* 42:811–814. <https://doi.org/10.1002/anie.200390216>
- Wang SX, Wang MX, Wang DX, Zhu J (2007) Chiral salen - aluminum complex as a catalyst for enantioselective  $\alpha$ -addition of isocyanides to aldehydes: asymmetric synthesis of 2-(1-hydroxyalkyl)-5-aminooxazoles. *Org Lett* 9:3615–3618. <https://doi.org/10.1021/ol7014658>
- Yue T, Wang MX, Wang DX, Masson G, Zhu J (2009) Catalytic asymmetric Passerini-type reaction: chiral aluminum-organophosphate-catalyzed enantioselective  $\alpha$ -addition of isocyanides to aldehydes. *J Org Chem* 74:8396–8399. <https://doi.org/10.1021/jo9017765>
- Yue T, Wang MX, Wang DX, Masson G, Zhu J (2009) Brønsted acid catalyzed enantioselective three-component reaction involving the  $\alpha$  addition of isocyanides to imines. *Angew Chem Int Ed* 48:6717–6721. <https://doi.org/10.1002/anie.200902385>
- Odabachian Y, Wang Q, Zhu J (2013) Tertiary amines as synthetic equivalents of vinyl cations: zinc bromide promoted coupling of propargylamines with  $\alpha$ -isocynoacetamides to give 2,4,5-trisubstituted oxazoles initiated by an internal redox

- process. *Chem Eur J* 19:12229–12233. <https://doi.org/10.1002/chem.201302106>
30. Mossetti R, Pirall T, Tron GC, Zhu J (2010) Efficient synthesis of  $\alpha$ -ketoamides via 2-acyl-5-aminooxazoles by reacting acyl chlorides and  $\alpha$ -isocyanoacetamides. *Org Lett* 12:820–823. <https://doi.org/10.1021/ol902894p>
  31. Janvier P, Sun X, Bienayme H, Zhu J (2002) Ammonium chloride-promoted four-component synthesis of pyrrolo[3,4-*b*]pyridin-5-one. *J Am Chem Soc* 124:2560–2567. <https://doi.org/10.1021/ja017563a>
  32. Janvier P, Bienaymé H, Zhu J (2002) A five-component synthesis of hexasubstituted benzene. *Angew Chem Int Ed* 41:4291–4294. [https://doi.org/10.1002/1521-3773\(20021115\)41:22%3c4291::AID-ANIE4291%3e3.0.CO;2-D](https://doi.org/10.1002/1521-3773(20021115)41:22%3c4291::AID-ANIE4291%3e3.0.CO;2-D)
  33. González-Zamora E, Fayol A, Bois-Choussy M, Chiaroni A, Zhu J (2001) Three component synthesis of oxa-bridged tetracyclic tetrahydroquinolines. *Chem Commun* 1:1684–1685. <https://doi.org/10.1039/b104317j>
  34. Fayol A, González-Zamora E, Bois-Choussy M, Zhu J (2007) Lithium bromide-promoted three-component synthesis of oxa-bridged tetracyclic tetrahydroisoquinolines. *Heterocycles* 73:729–742. [https://doi.org/10.3987/COM-07-S\(U\)55](https://doi.org/10.3987/COM-07-S(U)55)
  35. Montañó RG, Zhu J (2002) Rapid access to tetracyclic ring system of lennoxamine type natural product by combined use of a novel three-component reaction and Pummerer cyclization. *Chem Commun* 2002:2448–2449. <https://doi.org/10.1039/b207179g>
  36. Fayol A, Zhu J (2002) Synthesis of furoquinolines by a multi-component domino process. *Angew Chem Int Ed* 41:3633–3635. [https://doi.org/10.1002/1521-3773\(20021004\)41:19%3c3633::AID-ANIE3633%3e3.0.CO;2-T](https://doi.org/10.1002/1521-3773(20021004)41:19%3c3633::AID-ANIE3633%3e3.0.CO;2-T)
  37. Fayol A, Zhu J (2004) Synthesis of polysubstituted 4,5,6,7-tetrahydrofuro[2,3-*c*]pyridines by a novel multicomponent reaction. *Org Lett* 6:115–118. <https://doi.org/10.1021/ol036167p>
  38. Fayol A, Zhu J (2005) Multicomponent synthesis of epoxy-tetrahydronaphthridine and structural diversification by subsequent fragmentation. *Tetrahedron* 61:11511–11519. <https://doi.org/10.1016/j.tet.2005.09.017>
  39. Fayol A, Zhu J (2005) Three-component synthesis of polysubstituted 6-azaindolines and its tricyclic derivatives. *Org Lett* 7:239–242. <https://doi.org/10.1021/ol0477881>
  40. Bossio R, Marcaccini S, Pepino R (1990) Studies on isocyanides. Synthesis of *N*-substituted 2-isocyanocarboxamides. *Liebigs Ann Chem* 90:935–937. <https://doi.org/10.1002/jlac.1990199001172>
  41. Bossio R, Marcaccini S, Pepino R, Pellegrini G, Polsinelli M (1992) Synthesis of *N*-substituted isocyanocarboxamides with antimicrobial activity. *Farmaco* 47:1173–1179
  42. Bossio R, Marcaccini S, Pellegrini G, Pepino R (1992) *N*-substituted 2-isocyanoylacetamides with antimicrobial activity. *Arzneimittel-Forschung/Drug Research* 42–2:1494–1497
  43. Bossio R, Marcaccini S, Paoli P, Papaleo S, Pepino R, Polo C (1991) Studies on isocyanides and related compounds—synthesis and cyclization of *N*-substituted 1-isocyano-1- cycloalkanecarboxamides. *Liebigs Ann Chem* 1991:843–849. <https://doi.org/10.1002/jlac.1991199101145>
  44. Bossio R, Marcaccini S, Papaleo S, Pepino R (1994) Studies on isocyanides and related compounds - a convenient synthesis of 2,3-disubstituted spiroimidazolones. *J Heterocycl Chem* 31:397–399. <https://doi.org/10.1002/jhet.5570310225>
  45. Frippiat S, Leterrier C, Baudequin C, Hoarau C, Bischoff L (2020) Formation of imidazolones by ring closure of  $\alpha$ -isocyanoamides: Exploring new reactivities. *Synlett* 31:1211–1215. <https://doi.org/10.1055/s-0040-1708015>
  46. Muselli M, Baudequin C, Hoarau C, Bischoff L (2015) Pd-catalyzed direct C-H functionalization of imidazolones with aryl- and alkenyl halides. *Chem Commun* 51:745–748. <https://doi.org/10.1039/c4cc07917e>
  47. Gesù A, Pozzoli C, Torre E, Aprile S, Pirali T (2016) Transition-metal-free synthesis of 2-arylimidazolones via cascade reaction between arynes and  $\alpha$ ,  $\alpha'$ -disubstituted  $\alpha$ -isocyanoacetamides. *Org Lett* 18:1992–1995. <https://doi.org/10.1021/acs.orglett.6b00580>
  48. Pirali T, Tron GC, Masson G, Zhu J (2007) Ammonium chloride promoted three-component synthesis of 5-iminooxazoline and its subsequent transformation to macrocyclopeptide. *Org Lett* 9:5275–5278. <https://doi.org/10.1021/ol7024372>
  49. Djukanovic D, Petkovic M, Simic M, Jovanovic P, Tasic G, Savic V (2018) Synthesis of 2-unsubstituted imidazolones from bisamides via a one-pot, domino dehydration/base promoted cyclisation process. *Tetrahedron Lett* 59:914–917. <https://doi.org/10.1016/j.tetlet.2018.01.074>
  50. Waibel KA, Nickisch R, Möhl N, Seim R, Meier MAR (2020) A more sustainable and highly practicable synthesis of aliphatic isocyanides. *Green Chem* 22:933–941. <https://doi.org/10.1039/c9gc04070f>
  51. Bossio R, Marcaccini S, Pepino R, Polo C, Torroba T (1990) A facile synthesis of 1-aryl-2-arylthio-1*H*-imidazoles. *Heterocycles* 31:1287–1290. <https://doi.org/10.3987/COM-90-5390>
  52. Bossio R, Marcaccini S, Pepino R (1996) Studies on isocyanides and related compounds. Synthesis of 1-aryl-2-(tosylamino)-1*H*-imidazoles, a novel class of imidazole derivatives. *J Org Chem* 61:2202–2203. <https://doi.org/10.1021/jo9519165>
  53. Bossio R, Marcaccini S, Pepino R (1993) Studies on isocyanides and related compounds. A facile synthesis of imidazo[1,5-*a*]imidazoles. *Liebigs Ann Chem* 1993:1229–1231. <https://doi.org/10.1002/jlac.1993199301199>
  54. Patil P, Khoury K, Eberhardt H, Dömling A (2014) A universal isocyanide for diverse heterocycle synthesis. *Org Lett* 16:5736–5739. <https://doi.org/10.1021/ol5024882>
  55. Kazmaier U, Ackermann S (2005) A straightforward approach towards thiazoles and endotheiopeptides via Ugi reaction. *Org Biomol Chem* 3:3184–3187. <https://doi.org/10.1039/B507028G>
  56. Gulevich AV, Balenkova ES, Nenajdenko VG (2007) The first example of a diastereoselective thio-Ugi reaction: a new synthetic approach to chiral imidazole derivatives. *J Org Chem* 72:7878–7885. <https://doi.org/10.1021/jo071030o>
  57. Gunawan S, Hulme C (2013) Construction of functionalized tricyclic dihydropurazino-quinazolinedione chemotypes via an Ugi/*N*-acyliminium ion cyclization cascade. *Tetrahedron Lett* 54:4467–4470. <https://doi.org/10.1016/j.tetlet.2013.06.042>
  58. Meerwein H, Florian W, Schön N, Stopp G (1961) Über säureamidacetale, harnstoffacetale und lactamacetale. *Liebigs Ann Chem* 641:1–39. <https://doi.org/10.1002/jlac.19616410102>
  59. Dömling A, Illgen K (2005) 1-isocyano-2-dimethylamino-alkenes: versatile reagents in diversity-oriented organic synthesis. *Synthesis* 2005:662–667. <https://doi.org/10.1055/s-2004-831236>
  60. Schollkopf U, Porsch P-H, Lau H-H (1979) Notiz über *p*-dimethylamino- $\alpha$ -isocyanacrylsäureester und ihre verwendung in der heterocyclenchemie. *Liebigs Ann Chem* 1979:1444–1446. <https://doi.org/10.1002/jlac.197919790918>
  61. Lau H, Schollkopf U (1982) Syntheses with  $\alpha$ -metalated isocyanides. Synthesis of methyl 2-alkyl-1-methylimidazole-4-carboxylates and 2-acyl-1-methylimidazole-4-carboxylates from methyl (*Z*)- $\beta$ -dimethylamino- $\alpha$ -isocyanoacrylate and alkyl or acyl halides. *Liebigs Ann Chem* 11:2093–2095. <https://doi.org/10.1002/jlac.198219821117>
  62. Bossio R, Marcaccini S, Pepino R, Paoli P, Polo C (1993) Studies on isocyanides and related compounds—investigation on the reaction between (*Z*)-alkyl 3-dimethylamino-2- isocyanacrylates

- and acyl chlorides. *J Heterocycl Chem* 30:575–577. <https://doi.org/10.1002/jhet.5570300256>
63. Bossio R, Marcaccini S, Pepino R (1993) Studies on isocyanides and related compounds - a novel synthetic route to oxazole derivatives. *J Chem Res Synop* 1993:1
64. Bossio R, Marcaccini S, Pepino R, Paoli P (1994) Studies on isocyanides and related compounds - an unusual synthesis of imidazolylloxazolones. *J Heterocycl Chem* 31:729–732. <https://doi.org/10.1002/jhet.5570310406>
65. Helal CJ, Lucas JC (2002) A concise and regioselective synthesis of 1-alkyl-4-imidazolecarboxylates. *Org Lett* 4:4133–4134. <https://doi.org/10.1021/ol026892k>
66. Bienaymé H, Bouzid K (1998) Synthesis of rigid hydrophobic tetrazoles using an Ugi multi-component heterocyclic condensation. *Tetrahedron Lett* 39:2735–2738. [https://doi.org/10.1016/S0040-4039\(98\)00283-4](https://doi.org/10.1016/S0040-4039(98)00283-4)
67. Illgen K, Nerdinger S, Fuchs T, Friedrich C, Weber L, Herdtweck E (2004) A versatile synthesis of 6-oxo-1,4,5,6-tetrahydro-pyrazine-2-carboxylic acid methyl esters via MCR chemistry. *Synlett* 2004:53–56. <https://doi.org/10.1055/s-2003-43357>
68. Bienaymé H (1998) “Reagent explosion”: an efficient method to increase library size and diversity. *Tetrahedron Lett* 39:4255–4258. [https://doi.org/10.1016/S0040-4039\(98\)00696-0](https://doi.org/10.1016/S0040-4039(98)00696-0)
69. Heck S, Dömling A (2000) A versatile multi-component one-pot thiazole synthesis. *Synlett* 2000:424–426. <https://doi.org/10.1055/s-2000-6517>
70. Kolb J, Beck B, Almstetter M, Heck S, Herdtweck E, Dömling A (2003) New MCRs: the first 4-component reaction leading to 2,4-disubstituted thiazoles. *Mol Divers* 6:297–313. <https://doi.org/10.1023/B:MODI.0000006827.35029.e4>
71. Henkel B, Sax M, Dömling A (2003) A new and efficient multi-component solid-phase synthesis of 2-acylaminoethylthiazoles. *Tetrahedron Lett* 44:3679–3682. [https://doi.org/10.1016/S0040-4039\(03\)00662-2](https://doi.org/10.1016/S0040-4039(03)00662-2)
72. Zhang H (2014) A novel one-pot multicomponent enzymatic synthesis of 2,4-disubstituted thiazoles. *Catal Lett* 144:928–934. <https://doi.org/10.1007/s10562-014-1229-1>
73. Kolb J, Beck B, Dömling A (2002) Simultaneous assembly of the  $\beta$ -lactam and thiazole moiety by a new multicomponent reaction. *Tetrahedron Lett* 43:6897–6901. [https://doi.org/10.1016/S0040-4039\(02\)01621-0](https://doi.org/10.1016/S0040-4039(02)01621-0)
74. Wang W, Joyner S, Andrew K, Khoury S, Dömling A (2010) (-)-bacillamide C: the convergent approach. *Org Biomol Chem* 8:529–532. <https://doi.org/10.1039/b918214d>
75. Henkel B, Beck B, Westner B, Mejat B, Do A (2003) Convergent multicomponent assembly of 2-acyloxymethyl thiazoles. *Tetrahedron Lett* 44:8947–8950. <https://doi.org/10.1016/j.tetlet.2003.10.012>
76. Dömling A, Beck B, Eichelberger U, Sakamuri S, Menon S, Chen QZ, Lu Y, Wessjohann LA (2006) Total synthesis of tubulysin U and V. *Angew Chem Int Ed* 45:7235–7239. <https://doi.org/10.1002/anie.200601259>
77. Pando O, Stark S, Denkert A, Porzel A, Preusentanz R, Wessjohann LA (2011) The multiple multicomponent approach to natural product mimics: Tubugis, *N*-substituted anticancer peptides with picomolar activity. *J Am Chem Soc* 133:7692–7695. <https://doi.org/10.1021/ja2022027>
78. Bossio R, Marcaccini S, Pepino R, Torroba T (1996) Studies on isocyanides. 2-isocyanothioanisole, a synthetic equivalent of the benzothiazol-2-yl anion. *Heterocycles* 43:471–474. <https://doi.org/10.3987/COM-95-7318>
79. Yang W-C, Wei K, Sun X, Zhu J, Wu L (2018) Cascade C(sp<sup>3</sup>)-S bond cleavage and imidoyl C-S formation: radical cyclization of 2-isocyanophenyl thioethers toward 2-substituted benzothiazoles. *Org Lett* 20:3144–3147. <https://doi.org/10.1021/acs.orglett.8b01278>
80. Liu Y, Li J-L, Liu X-G, Wu J-Q, Huang Z-S, Li Q, Wang H (2021) Radical borylative cyclization of isocyanophenyls with *N*-heterocyclic carbene borane: synthesis of borylated azarenes. *Org Lett* 23:1891–1897. <https://doi.org/10.1021/acs.orglett.1c00309>
81. Luo K, Yang W-C, Wei K, Liu Y, Wang J-K, Wu L (2019) Di-*tert*-butyl peroxide-mediated radical C(sp<sup>2</sup>/sp<sup>3</sup>)-S bond cleavage and group-transfer cyclization. *Org Lett* 21:7851–7856. <https://doi.org/10.1021/acs.orglett.9b02837>
82. Yuan Y, Dong W, Gao X, Xie X, Zhang Z (2019) Sodium sulfite-involved photocatalytic radical cascade cyclization of 2-isocyanophenyl thioethers: Access to 2-CF<sub>2</sub>/CF<sub>3</sub>-containing benzothiazoles. *Org Lett* 21:469–472. <https://doi.org/10.1021/acs.orglett.8b03710>
83. Liu Y, Chen X-L, Sun K, Li X-Y, Zeng F-L, Liu X-C, Qu L-B, Zhao Y-F, Yu B (2019) Visible-light induced radical perfluoroalkylation/cyclization strategy to access 2-perfluoroalkylbenzothiazoles/benzoselenazoles by EDA complex. *Org Lett* 21:4019–4024. <https://doi.org/10.1021/acs.orglett.9b01175>
84. Xie XY, Li Y, Xia YT, Luo K, Wu L (2021) Visible light-induced metal-free and oxidant-free radical cyclization of (2-isocyanophenyl)(methyl)sulfanes with ethers. *Eur J Org Chem* 2021:4273–4277. <https://doi.org/10.1002/ejoc.202100501>
85. Liu Y, Chen X-L, Li X-Y, Zhu S-S, Li S-J, Song Y, Qu L-B, Yu B (2021) 4CzIPN-<sup>t</sup>Bu-catalyzed proton-coupled electron transfer for photosynthesis of phosphorylated *N*-heteroaromatics. *J Am Chem Soc* 143:964–972. <https://doi.org/10.1021/jacs.0c11138>
86. Banfi L, Basso A, Lambruschini C, Moni L, Riva R (2017) Synthesis of seven-membered nitrogen heterocycles through the Ugi multicomponent reaction. *Chem Heterocycl Compd* 53:382–408. <https://doi.org/10.1007/s10593-017-2065-1>
87. Hulme C, Dietrich J (2009) Emerging molecular diversity from the intra-molecular Ugi reaction: Iterative efficiency in medicinal chemistry. *Mol Divers* 13:195–207. <https://doi.org/10.1007/s11030-009-9111-6>
88. Short KM, Ching BW, Mjalli AMM (1997) Exploitation of the Ugi 4CC reaction: preparation of small molecule combinatorial libraries via solid phase. *Tetrahedron* 53:6653–6679. [https://doi.org/10.1016/S0040-4020\(97\)00223-8](https://doi.org/10.1016/S0040-4020(97)00223-8)
89. Harriman GCB (1997) Synthesis of small and medium sized 2,2-disubstituted lactams via the “intramolecular” three component Ugi reaction. *Tetrahedron Lett* 38:5591–5594. [https://doi.org/10.1016/S0040-4039\(97\)01265-3](https://doi.org/10.1016/S0040-4039(97)01265-3)
90. Hanusch-Kompa C, Ugi I (1998) Multi-component reactions 13: Synthesis of  $\gamma$ -lactams as part of a multiring system via Ugi-4-centre-3-component reaction. *Tetrahedron Lett* 39:2725–2728. [https://doi.org/10.1016/S0040-4039\(98\)00428-6](https://doi.org/10.1016/S0040-4039(98)00428-6)
91. Marcaccini S, Pepino R, Torroba T, Miguel D, Garcia-Valverde M (2002) Synthesis of thiomorpholines by an intramolecular Ugi reaction. *Tetrahedron Lett* 43:8591–8593. [https://doi.org/10.1016/S0040-4039\(02\)02064-6](https://doi.org/10.1016/S0040-4039(02)02064-6)
92. Marcaccini S, Miguel D, Torroba T, Garcia-Valverde M (2003) 1,4-thiazepines, 1,4-benzothiazepin-5-ones, and 1,4-benzothioxepin orthoamides via multicomponent reactions of isocyanides. *J Org Chem* 68:3315–3318. <https://doi.org/10.1021/jo026614z>
93. Ilyn AP, Loseva MV, Vvedensky VY, Putsykina EB, Tkachenko SE, Kravchenko DV, Khvat AV, Krasavin MY, Ivachtchenko AV (2006) One-step assembly of carbamoyl-substituted heteroannulated [1,4]thiazepines. *J Org Chem* 71:2811–2819. <https://doi.org/10.1021/jo052640w>
94. Ilyn AP, Parchinski VZ, Peregudova JN, Trifilenkov AS, Putsykina EB, Tkachenko SE, Kravchenko DV, Ivachtchenko AV (2006) One-step assembly of carbamoyl substituted annulated 1,4-oxazepines. *Tetrahedron Lett* 47:2649–2653

95. Potapov VV, Fetisova NA, Nikitina AV, Ivachtchenko AV (2009) A convenient synthesis of heterocyclic compounds containing 11-oxo-6,11,12,13-tetrahydrodibenzo[*b*, *g*][1,5]oxazonine fragment. *Mendeleev Commun* 19:287–289. <https://doi.org/10.1016/j.mencom.2009.09.020>
96. Zhang J, Jacobson A, Rusche JR, Herlihy W (1999) Unique structures generated by Ugi 3CC reactions using bifunctional starting materials containing aldehyde and carboxylic acid. *J Org Chem* 64:1074–1076. <https://doi.org/10.1021/jo982192a>
97. Krasavin M, Parchinsky V, Shumsky A, Konstantinov I, Vantskul A (2010) Proline-like  $\beta$ -turn mimics accessed via Ugi reaction involving monoprotected hydrazines. *Tetrahedron Lett* 51:1367–1370. <https://doi.org/10.1016/j.tetlet.2009.12.141>
98. Cioc RC, Estévez V, van der Niet DJ, Vande Velde CML, Turrini NG, Hall M, Faber K, Ruijter E, Orru RVA (2017) Stereoselective synthesis of functionalized bicyclic scaffolds by Passerini 3-center-2-component reactions of cyclic ketoacids. *Eur J Org Chem* 17:1262–1271. <https://doi.org/10.1002/ejoc.201601432>
99. Ley SV, Taylor SJ (2002) A polymer-supported [1,3,2] oxazaphospholidine for the conversion of isothiocyanates to isocyanides and their subsequent use in an Ugi reaction. *Bioorg Med Chem Lett* 12:1813–1816. [https://doi.org/10.1016/s0960-894x\(02\)00269-x](https://doi.org/10.1016/s0960-894x(02)00269-x)
100. Faggi C, García-Valverde M, Marcaccini S, Menchi G (2010) Isolation of Ugi four-component condensation primary adducts: a straightforward route to isocoumarins. *Org Lett* 12:788–791. <https://doi.org/10.1021/ol9028622>
101. Ramazani A, Mahyari A (2010) Three-component reaction of isocyanides and 2-formylbenzoic acid with dibenzylamine catalyzed by silica nanoparticles under solvent-free conditions. *Helv Chim Acta* 93:2203–2209. <https://doi.org/10.1002/hlca.201000124>
102. Naghdipari K, Shojaei L, Heidari A, Heidarifard M, Sharbati M, Mahari A, Hosseinzadeh-Khanmiri R (2019) Isochromone-functionalized mesoporous silica hollow sphere as an efficient material for drug delivery. *Polyhedron* 170:659–665. <https://doi.org/10.1016/j.poly.2019.06.035>
103. Marcaccini S, Menchi G, Trabocchi A (2011) Synthesis of diverse phenylglycine derivatives via transformation of Ugi four-component condensation primary adducts. *Tetrahedron Lett* 52:2673–2675. <https://doi.org/10.1016/j.tetlet.2011.03.074>
104. Zhang Y, Ao Y-F, Huang Z-T, Wang D-X, Wang M-X, Zhu J (2016) Chiral phosphoric acid catalyzed asymmetric Ugi reaction by dynamic kinetic resolution of the primary multicomponent adduct. *Angew Chem Int Ed* 55:5282–5285
105. Esmaili-Marandi F, Yavari I, Saeedi M, Mahdavi M, Shafiee A (2016) Synthesis of novel pyrazino[2,1-*a*]isoindolodiones via intramolecular hydroamination of 2,3-dihydro-3-oxo-2-(prop-2-yn-1-yl)-1*H*-isoindole-1-carboxamides. *Helv Chim Acta* 99:187–190
106. Bachman M, Mann SE, Sheppard TD (2012) Rapid synthesis of highly functionalised  $\alpha$ -amino amides and medium ring lactones using multicomponent reactions of amino alcohols and isocyanides. *Org Biomol Chem* 10:162–170. <https://doi.org/10.1039/C1OB06534C>
107. Ma G-H, Jiang B, Tu X-J, Ning Y, Tu S-J, Li G (2014) Synthesis of isocoumarins with different substituted patterns via Passerini-aldol sequence. *Org Lett* 16:4504–4507. <https://doi.org/10.1021/ol502048e>
108. Ponra S, Nyadanu A, Kaïm LE, Grimaud L, Vitale MR (2016)  $\text{TiCl}_4$ -mediated preparation of thiophthalide derivatives via formal thio-Passerini reactions. *Org Lett* 18:4060–4063. <https://doi.org/10.1021/acs.orglett.6b01937>
109. Vitale M, Grimaud L, El Kaïm L, Ponra S, Nyadanu A, Maurin S (2017)  $\text{TiCl}_4$ -mediated synthesis of 3,4-hetero-disubstituted isocoumarins by means of isocyanide insertion reactions. *Synthesis* 50:1331–1342. <https://doi.org/10.1055/s-0036-1591733>
110. Bossio R, Marcaccini S, Paoli P, Pepino R (1994) Synthesis of *N*, *N'*-diaryldipeptide amides by multicomponent condensation utilizing potassium isocynoacetate. *Synthesis* 1994:672–674. <https://doi.org/10.1055/s-1994-25541>
111. Bonne D, Dekhane M, Zhu J (2004) Ammonium chloride promoted Ugi four-component, five-center reaction of  $\alpha$ -substituted  $\alpha$ -isocyano acetic acid: a strong solvent effect. *Org Lett* 6:4771–4774. <https://doi.org/10.1021/ol0479388>
112. Bonne D, Dekhane M, Zhu J (2005) Mild oxidative one-carbon homologation of aldehyde to amide. *J Am Chem Soc* 127:6926–6927. <https://doi.org/10.1021/ja0511220>
113. Bossio R, Marcos CF, Marcaccini S, Pepino R (1997) A facile synthesis of  $\beta$ -lactams based on the isocyanide chemistry. *Tetrahedron Lett* 38:2519–2520. [https://doi.org/10.1016/s0040-4039\(97\)00389-4](https://doi.org/10.1016/s0040-4039(97)00389-4)
114. Stevens CV, Vekemans W, Moonen K, Rammeloo T (2003) Synthesis of 4-phosphono- $\beta$ -lactams via phosphite addition to acyliminium salts. *Tetrahedron Lett* 44:1619–1622. [https://doi.org/10.1016/S0040-4039\(03\)00005-4](https://doi.org/10.1016/S0040-4039(03)00005-4)
115. Moonen K, Stevens CV (2005) One-pot synthesis of *N*-chloroacetyl 1-aminoalkyl phosphonates - precursors of 4-phosphono- $\beta$ -lactams. *Synthesis* 2005:3603–3612. <https://doi.org/10.1055/s-2005-918440>
116. Van Speybroeck V, Moonen K, Hemelsoet K, Stevens CV, Waroquier M (2006) Unexpected four-membered over six-membered ring formation during the synthesis of azaheterocyclic phosphonates: Experimental and theoretical evaluation. *J Am Chem Soc* 128:8468–8478. <https://doi.org/10.1021/ja0584119>
117. Gerona-Navarro G, Bonache MA, Herranz R, García-López MT, González-Muñiz R (2001) Entry to new conformationally constrained amino acids. First synthesis of 3-unsubstituted 4-alkyl-4-carboxy-2-azetidinone derivatives via an intramolecular  $\text{N}^\alpha\text{-C}^\alpha$ -cyclization strategy. *J Org Chem* 66:3538–3547. <https://doi.org/10.1021/jo015559b>
118. Bonache MA, Gerona-Navarro G, Martín-Martínez M, García-López MT, López P, Cativiela C, González-Muñiz R (2003) Memory of chirality in the enantioselective synthesis of  $\beta$ -lactams derived from amino acids Influence of the reaction conditions. *Synlett* 2003:1007–1011. <https://doi.org/10.1055/s-2003-39301>
119. Bossio R, Marcos CF, Marcaccini S, Pepino R (1997) Studies on isocyanides and related compounds. An unusual synthesis of functionalized succinimides. *Synthesis* 1997:1389–1390. <https://doi.org/10.1055/s-1997-1367>
120. Bossio R, Marcaccini S, Pepino R, Marcos CF (2000) Multicomponent reactions - a convenient undergraduate organic chemistry experiment. *J Chem Educ* 77:382–384. <https://doi.org/10.1021/ed077p382>
121. Ghabraie E, Balalaie S, Mehrparvar S, Rominger F (2014) Synthesis of functionalized  $\beta$ -lactams and pyrrolidine-2,5-diones through a metal-free sequential Ugi-4CR/cyclization reaction. *J Org Chem* 79:7926–7934. <https://doi.org/10.1021/jo5010422>
122. Yan Y-M, Wang M-L, Liu Y-L, He Y-C (2020) One-pot and regioselective synthesis of functionalized  $\gamma$ -lactams via a metal-free sequential Ugi 4CR/intramolecular 5-*exo-dig* cyclization reaction. *Tetrahedron* 76:131389–131396. <https://doi.org/10.1016/j.tet.2020.131389>
123. Bossio R, Marcos CF, Marcaccini S, Pepino R (1997) Studies on isocyanides and related compounds. A novel synthetic route to 1,6-dihydro-6-oxopyridine-2-carboxylic acid derivatives. *Heterocycles* 45:1589–1592. <https://doi.org/10.3987/COM-97-7855>
124. Bossio R, Marcaccini S, Pepino R (1994) Studies on isocyanides and related compounds - a novel synthesis of pyrroles

- via Ugi reaction. *Synthesis* 1994:765–766. <https://doi.org/10.1055/s-1994-25565>
125. Marcos CF, Marcaccini S, Pepino R, Polo C, Torroba T (2003) Studies on isocyanides and related compounds; a facile synthesis of functionalized 3(2*H*)-pyridazinones via Ugi four-component condensation. *Synthesis* 2003:691–694. <https://doi.org/10.1055/s-2003-38071>
126. Marcaccini S, Pepino R, Pozo MC, Basurto S, Maria GVB, Torroba T (2004) One-pot synthesis of quinolin-2-(1*H*)-ones via tandem Ugi-Knoevenagel condensations. *Tetrahedron Lett* 45:3999–4001. <https://doi.org/10.1016/j.tetlet.2004.03.184>
127. Neo AG, Carrillo RM, Barriga S, Moman E, Marcaccini S, Marcos CF (2007) Multicomponent synthesis of highly substituted 2-pyridones. *Synlett* 2007:327–329. <https://doi.org/10.1055/s-2007-967992>
128. Alvarez-Rodríguez NV, Dos Santos A, El Kaïm L, Gámez-Montaño R (2015) The Ugi reaction of cyanoacetic acid as a route to tetramic acid derivatives. *Synlett* 26:2253–2256. <https://doi.org/10.1055/s-0035-1560050>
129. Trifilenkov AS, Ilyin AP, Kysil VM, Sandulenko YB, Ivachtchenko AV (2007) One-pot tandem complexity-generating reaction based on Ugi four component condensation and intramolecular cyclization. *Tetrahedron Lett* 48:2563–2567. <https://doi.org/10.1016/j.tetlet.2007.02.015>
130. Zhang L, Zhao F, Zheng M, Zhai Y, Liu H (2013) Rapid and selective access to three distinct sets of indole-based heterocycles from a single set of Ugi-adducts under microwave heating. *Chem Commun* 49:2894. <https://doi.org/10.1039/c3cc00111c>
131. Polindara-García LA, Miranda LD (2012) Two-step synthesis of 2,3-dihydropyrroles via a formal 5-endo cycloisomerization of Ugi 4-CR/propargyl adducts. *Org Lett* 14:5408–5411. <https://doi.org/10.1021/ol3024727>
132. Polindara-García LA, Vazquez A (2014) Combinatorial synthesis of nicotine analogs using an Ugi 4-CR/cyclization-reduction strategy. *Org Biomol Chem* 12:7068. <https://doi.org/10.1039/c4ob00767k>
133. Shahriari A, Amiri K, Nikbakht A, Rominger F, Bijanzadeh HR, Balalaie S (2022) Synthesis of pyrrolidin-5-one-2-carboxamides through cyclization of *N*-substituted-2-alleneamides. *J Org Chem* 87:7778–7785. <https://doi.org/10.1021/acs.joc.2c00387>
134. Polindara-García LA, Montesinos-Miguel D, Vazquez A (2015) An efficient microwave-assisted synthesis of cotinine and isocotinine analogs from an Ugi-4CR approach. *Org Biomol Chem* 13:9065–9071. <https://doi.org/10.1039/c5ob01170a>
135. Shiri M, Mirpour-Marzoni SZ, Bozorgpour-Savadjani Z, Soleymanifard B, Kruger HG (2014) Base-catalyzed cyclization of Ugi-adducts to substituted indolyl based  $\gamma$ -lactams. *Monatsh Chem* 145:1947–1952. <https://doi.org/10.1007/s00706-014-1271-0>
136. Che C, Li S, Jiang X, Quan J, Lin S, Yang Z (2010) One-pot syntheses of chromeno[3,4-*c*]pyrrole-3,4-diones via Ugi-4CR and intramolecular Michael addition. *Org Lett* 12:4682–4685. <https://doi.org/10.1021/ol1020477>
137. Ben Abdessalem A, Abderrahim R, Agrebie A, Dos Santos A, El Kaïm L, Komesky A (2015) Formal [3+2] cycloaddition of Ugi adducts towards pyrrolines. *Chem Commun* 51:1116–1119. <https://doi.org/10.1039/c4cc08282f>
138. Jia S, El Kaïm L (2018) A Passerini/Michael pathway towards butyrolactones. *Eur J Org Chem* 2018:6457–6464. <https://doi.org/10.1002/ejoc.201800958>
139. Zidan A, Garrec J, Cordier M, El-Naggar AM, Abd El-Sattar NEA, Ali AK, Hassan MA, El Kaim L (2017) *B*-lactam synthesis through diodomethane addition to amide dianions. *Angew Chem Int Ed* 56:12179–12183. <https://doi.org/10.1002/anie.201706315>
140. Zidan A, Cordier M, El-Naggar AM, Abd El-Sattar NEA, Hassan MA, Ali AK, El Kaïm L (2018) Propargylation of Ugi amide dianion: an entry into pyrrolidinone and benzoindolizidine alkaloid analogues. *Org Lett* 20:2568–2571. <https://doi.org/10.1021/acs.orglett.8b00687>
141. Salcedo A, Neuville L, Zhu J (2008) Palladium-catalyzed intramolecular *C*-arylation of benzylic carbon: synthesis of 3-benzoxazolyloindolinones by a sequence of Ugi-4CR/postfunctionalization. *J Org Chem* 73:3600–3603. <https://doi.org/10.1021/jo800266y>
142. Ghandi M, Zarezadeh N, Abbasi A (2015) One-pot synthesis of spiropyrroloquinoline-isoindolinone and their aza-analogs via the Ugi-4CR/metal-free intramolecular bis-annulation process. *Org Biomol Chem* 13:8211–8220. <https://doi.org/10.1039/c5ob01095k>
143. Butani HH, Vachhani DD, Bhoya UC, Shah AK, Van Der Eycken EV (2014) Regio- and chemoselective formation of spiroindolinone-isoindolinone by a palladium-catalyzed Buchwald-Hartwig addition-elimination sequence. *Eur J Org Chem* 2014:6634–6638. <https://doi.org/10.1002/ejoc.201402909>
144. Sharma N, Li Z, Sharma UK, Van Der Eycken EV (2014) Facile access to functionalized spiro[indoline-3,2'-pyrrole]-2,5'-diones via post-Ugi domino Buchwald-Hartwig/Michael reaction. *Org Lett* 16:3884–3887. <https://doi.org/10.1021/ol5019079>
145. El Kaïm L, Grimaud L, Le Goff X-F, Menes-Arzate M, Miranda LD (2011) Straightforward four-component access to spiroindolines. *Chem Commun* 47:8145. <https://doi.org/10.1039/c1cc12236c>
146. Zidan A, El-Naggar AM, Abd El-Sattar NEA, Ali AK, El Kaim L (2019) Raising the diversity of Ugi reactions through selective alkylations and allylations of Ugi adducts. *Front Chem* 7:20. <https://doi.org/10.3389/fchem.2019.00020>
147. Tyagi V, Khan S, Chauhan PMS (2013) Facile synthesis of diverse isoindolinone derivatives via Ugi-4CR followed by Cu-catalyzed deamidative C(sp<sup>2</sup>)-C(sp<sup>3</sup>) coupling. *Tetrahedron Lett* 54:1279–1284. <https://doi.org/10.1016/j.tetlet.2012.12.091>
148. Trang TTT, Peshkov AA, Jacobs J, Van Meervelt L, Peshkov VA, Van der Eycken EV (2015) Post-Ugi carbocyclization/fragmentation sequence for the synthesis of 6,7-dihydro-5*H*-pyrrolo[3,4-*b*]pyridin-5-ones. *Tetrahedron Lett* 56:2882–2886. <https://doi.org/10.1016/j.tetlet.2015.03.092>
149. Ramiro JL, Neo AG, Marcos CF (2022) Synthesis of imidazolo-coumarins by the amide-directed oxidative cyclisation of enol-Ugi derivatives. *Org Biomol Chem* 20:5293–5307. <https://doi.org/10.1039/d2ob00518b>
150. Garcia-Valverde M, Macho S, Marcaccini S, Rodriguez T, Rojo J, Torroba T (2008) A one-pot, Ugi four-component synthesis of 2(3*H*)-oxazolone 4-carboxamides. *Synlett* 2008:33–36. <https://doi.org/10.1055/s-2007-992372>
151. Zeng X-H, Wang H-M, Yan Y-M, Wu L, Ding M-W (2014) One-pot regioselective synthesis of  $\beta$ -lactams by a tandem Ugi 4CC/S<sub>N</sub> cyclization. *Tetrahedron* 70:3647–3652. <https://doi.org/10.1016/j.tet.2014.04.033>
152. Zeng X-H, Wang H-M, Wu L, Ding M-W (2013) One-pot synthesis of 5-oxopyrrolidine-2-carboxamides via a tandem Ugi 4CC/S<sub>N</sub> cyclization starting from Baylis-Hillman bromides. *Tetrahedron* 69:3823–3828. <https://doi.org/10.1016/j.tet.2013.03.058>
153. Pertejo P, Sancho-Medina A, Hermosilla T, González-Saiz B, Gómez-Ayuso J, Quesada R, Moreno D, Carreira-Barral I, García-Valverde M (2021) Keto-enol tautomerism in Passerini and Ugi adducts. *Molecules* 26:919. <https://doi.org/10.3390/molecules26040919>
154. González-Saiz B, Pertejo P, Peña-Calleja P, Mielczarek M, Hermosilla T, Carreira-Barral I, De Miguel O, Rodríguez-Vidal F, Quesada R, García-Valverde M (2022) Base-selective access to highly functionalized heterocycles from multicomponent Ugi

- adducts. *Green Chem* 24:7988–7995. <https://doi.org/10.1039/d2gc02896d>
155. Peshkov AA, Peshkov VA, Li Z, Pereshivko OP, Van Der Eycken EV (2014) Assembly of a 1*H*-pyrrol-2(5*H*)-one core through a cascade Ugi reaction/5-*endo-dig* carbocyclization/retro-Claisen fragmentation process. *Eur J Org Chem* 2014:6390–6393. <https://doi.org/10.1002/ejoc.201402723>
156. Li Z, Kumar A, Peshkov A, Van Der Eycken EV (2016) A domino Ugi/Michael approach for the synthesis of  $\alpha$ ,  $\beta$ -unsaturated  $\gamma$ -lactams. *Tetrahedron Lett* 57:754–756. <https://doi.org/10.1016/j.tetlet.2016.01.014>
157. Van der Eycken E, Peshkov V, Nechaev A, Peshkov A (2016) The application of multicomponent Ugi and Passerini reactions for the one-pot synthesis of pyrrolones and butenolides. *Synthesis* 48:2280–2286. <https://doi.org/10.1055/s-0035-1561422>
158. Pertejo P, González-Saiz B, Quesada R, García-Valverde M (2020) One-pot synthesis of enantiopure pyrrolpiperazines. *J Org Chem* 85:14240–14245. <https://doi.org/10.1021/acs.joc.0c02103>
159. Marcaccini S, Pepino R, Pozo MC (2001) A facile synthesis of 2,5-diketopiperazines based on isocyanide chemistry. *Tetrahedron Lett* 42:2727–2728. [https://doi.org/10.1016/S0040-4039\(01\)00232-5](https://doi.org/10.1016/S0040-4039(01)00232-5)
160. Bruttomesso AC, Eiras J, Ramírez JA, Galagovsky LR (2009) Highly stereoselective synthesis of steroidal 2,5-diketopiperazines based on isocyanide chemistry. *Tetrahedron Lett* 50:4022–4024. <https://doi.org/10.1016/j.tetlet.2009.03.219>
161. Caputo S, Banfi L, Basso A, Galatini A, Moni L, Riva R, Lambruschini C (2017) Diversity-oriented synthesis of various enantiopure heterocycles by coupling organocatalysis with multicomponent reactions. *Eur J Org Chem* 2017:6619–6628. <https://doi.org/10.1002/ejoc.201701328>
162. Malet G, Martín AG, Orzáez M, Vicent MJ, Masip I, Sanclimens G, Ferrer-Montiel A, Mingarro I, Messeguer A, Fearnhead HO, Pérez-Payá E (2006) Small molecule inhibitors of apaf-1-related caspase-3/-9 activation that control mitochondrial-dependent apoptosis. *Cell Death Differ* 13:1523–1532. <https://doi.org/10.1038/sj.cdd.4401828>
163. Corredor M, Bujons J, Orzaez M, Sancho M, Perez-Paya E, Alfonso I, Messeguer A (2013) Optimizing the control of apoptosis by amide/triazole isosteric substitution in a constrained peptoid. *Eur J Med Chem* 63:892–896. <https://doi.org/10.1016/j.ejmech.2013.03.004>
164. Corredor M, Garrido M, Bujons J, Orzáez M, Pérez-Payá E, Alfonso I, Messeguer A (2015) Efficient synthesis of conformationally restricted apoptosis inhibitors bearing a triazole moiety. *Chem Eur J* 21:14122–14128. <https://doi.org/10.1002/chem.201502380>
165. Garrido M, Corredor M, Orzáez M, Alfonso I, Messeguer A (2016) Regioselective synthesis of a family of  $\beta$ -lactams bearing a triazole moiety as potential apoptosis inhibitors. *ChemistryOpen* 5:485–494. <https://doi.org/10.1002/open.201600052>
166. Vroemans R, Bamba F, Winters J, Thomas J, Jacobs J, Van Meervelt L, John J, Dehaen W (2018) Sequential Ugi reaction/base-induced ring closing/IAAC protocol toward triazolobenzodiazepine-fused diketopiperazines and hydantoins. *Beilstein J Org Chem* 14:626–633. <https://doi.org/10.3762/bjoc.14.49>
167. Hernández-Vázquez E, Miranda LD (2016) Practical synthesis and cytotoxic evaluation of the pyrazino[1,2-*b*]-isoquinoline ring system. *Org Biomol Chem* 14:4875–4884. <https://doi.org/10.1039/c6ob00431h>
168. Golubev P, Krasavin M (2017) Sterically constrained and encumbered: An approach to the naturally occurring peptidomimetic tetrahydropyrazino[1,2-*a*]indole-1,4-dione core. *Eur J Org Chem* 2017:1740–1744. <https://doi.org/10.1002/ejoc.201700152>
169. González-Saiz B, Carreira-Barral I, Pertejo P, Gómez-Ayuso J, Quesada R, García-Valverde M (2022) One-pot diastereoselective synthesis of pyrrolpiperazine-2,6-diones by a Ugi/nucleophilic substitution/*N*-acylation sequence. *J Org Chem* 87:9391–9398. <https://doi.org/10.1021/acs.joc.2c00694>
170. Ignacio JM, Macho S, Marcaccini S, Pepino R, Torroba T (2005) A facile synthesis of 1,3,5-trisubstituted hydantoins via Ugi four-component condensation. *Synlett* 2005:3051–3054. <https://doi.org/10.1055/s-2005-922745>
171. Hulme C, Ma L, Romano JJ, Morton G, Tang S-Y, Cherrier M-P, Choi S, Salvino J, Labaudiniere R (2000) Novel applications of carbon dioxide/MeOH for the synthesis of hydantoins and cyclic ureas via the Ugi reaction. *Tetrahedron Lett* 41:1889–1893. [https://doi.org/10.1016/S0040-4039\(00\)00053-8](https://doi.org/10.1016/S0040-4039(00)00053-8)
172. Chen Z-Z, Xu C, Xu Z-G, Ding Y, Meng J-P, Tang D-Y, Li Y, Lei J (2018) Facile construction of hydantoin scaffolds via a post-Ugi cascade reaction. *Synlett* 29:2199–2202. <https://doi.org/10.1055/s-0037-1610234>
173. Sañudo M, García-Valverde M, Marcaccini S, Torroba T (2012) A diastereoselective synthesis of pseudopeptidic hydantoins by an Ugi/cyclization/Ugi sequence. *Tetrahedron* 68:2621–2629. <https://doi.org/10.1016/j.tet.2012.01.073>
174. Brockmeyer F, Kroeger D, Stalling T, Ullrich P, Martens J (2012) A manifold three-step synthetic route to polycyclic annulated hydantoins via cyclic imines. *Helv Chim Acta* 95:1857–1870. <https://doi.org/10.1002/hlca.201200441>
175. Firth J, Zhang R, Morgentin R, Guilleux R, Kalliokoski T, Wariner S, Foster R, Marsden S, Nelson A (2015) Exploitation of the Ugi-Joullié reaction in the synthesis of libraries of drug-like bicyclic hydantoins. *Synthesis* 47:2391–2406. <https://doi.org/10.1055/s-0034-1378704>
176. Yu S-J, Zhu C, Bian Q, Cui C, Du X-J, Li Z-M, Zhao W-G (2014) Novel ultrasound-promoted parallel synthesis of trifluoroatrolactamide library via a one-pot Passerini/hydrolysis reaction sequence and their fungicidal activities. *ACS Comb Sci* 16:17–23. <https://doi.org/10.1021/co400111r>
177. Faggi C, Neo AG, Marcaccini S, Menchi G, Revuelta J (2008) Ugi four-component condensation with two cleavable components: the easiest synthesis of 2 *N*-diarylglycines. *Tetrahedron Lett* 49:2099–2102. <https://doi.org/10.1016/j.tetlet.2008.01.134>
178. Marcos CF, Marcaccini S, Menchi G, Pepino R, Torroba T (2008) Studies on isocyanides: Synthesis of tetrazolyl-isoindolinones via tandem Ugi four-component condensation/intramolecular amidation. *Tetrahedron Lett* 49:149–152. <https://doi.org/10.1016/j.tetlet.2007.10.154>
179. Nixey T, Kelly M, Hulme C (2000) The one-pot solution phase preparation of fused tetrazole-ketopiperazines. *Tetrahedron Lett* 41:8729–8733. [https://doi.org/10.1016/S0040-4039\(00\)01563-x](https://doi.org/10.1016/S0040-4039(00)01563-x)
180. Borisov RS, Polyakov AI, Medvedeva LA, Guranova NI, Voskressensky LG (2013) Synthesis of tetrazolodiazepines by a five-centered four-component azide Ugi reaction Scope and limitations. *Russ Chem Bull* 61:1609–1615. <https://doi.org/10.1007/s1172-012-0214-3>
181. Yerande SG, Newase KM, Singh B, Boltjes A, Dömling A (2014) Application of cyclic ketones in MCR: Ugi/amide coupling based synthesis of fused tetrazolo[1,5-*a*][1,4]benzodiazepines. *Tetrahedron Lett* 55:3263–3266. <https://doi.org/10.1016/j.tetlet.2014.04.040>
182. Boltjes A, Liao GP, Zhao T, Herdtweck E, Dömling A (2014) Ugi 4-CR synthesis of  $\gamma$ - and  $\delta$ -lactams providing new access to diverse enzyme interactions, a PDB analysis. *MedChemComm* 5:949–952. <https://doi.org/10.1039/C4MD00162A>
183. Bossio R, Marcaccini S, Paoli P, Pepino R, Polo C (1991) Studies on isocyanides and related compounds - synthesis of benzofuran derivatives. *Synthesis* 1991:999–1000. <https://doi.org/10.1055/s-1991-26628>



184. Dömling A (2006) Recent developments in isocyanide based multicomponent reactions in applied chemistry. *Chem Rev* 106:17–89. <https://doi.org/10.1021/cr0505728>
185. Ayaz M, De Moliner F, Dietrich J, Hulme C (2012) Applications of isocyanides in IMCRs for the rapid generation of molecular diversity. *Isocyanide chemistry*. <https://doi.org/10.1002/9783527652532.ch10>
186. Marcaccini S, Pepino R, Marcos CF, Polo C, Torroba T (2000) Studies on isocyanides and related compounds. Synthesis of furan derivatives and their transformation into indole derivatives. *J Heterocycl Chem* 37:1501–1503. <https://doi.org/10.1002/jhet.5570370615>
187. Faggi C, Marcaccini S, Pepino R, Pozo MC (2002) Studies on isocyanides and related compounds; synthesis of 1,4-benzodiazepine-2,5-diones via Ugi four-component condensation. *Synthesis* 2002:2756–2760. <https://doi.org/10.1055/s-2002-35987>
188. Marcaccini S, Miliciani M, Pepino R (2005) A facile synthesis of 1,4-benzodiazepine derivatives via Ugi four-component condensation. *Tetrahedron Lett* 46:711–713. <https://doi.org/10.1016/j.tetlet.2004.11.083>
189. Pertejo P, Garcia-Valverde M, Pena P, Cordero NA, Torroba T, Gonzalez-Ortega A (2014) Experimental and theoretical studies on the effect of the oxo group in 1,4-benzodiazepines. *Org Biomol Chem* 12:4905–4916. <https://doi.org/10.1039/c4ob00444b>
190. Guo Z, Zhuang C, Zhu L, Zhang Y, Yao J, Dong G, Wang S, Liu Y, Chen H, Sheng C, Miao Z, Zhang W (2012) Structure–activity relationship and antitumor activity of thio-benzodiazepines as p53–mdm2 protein–protein interaction inhibitors. *Eur J Med Chem* 56:10–16. <https://doi.org/10.1016/j.ejmech.2012.08.003>
191. Pertejo P, Pena-Calleja P, Carreira-Barral I, Quesada R, Alejandro Cordero N, Javier Rodriguez F, Garcia-Valverde M (2017) Novel pyrrolobenzodiazepine and pyrroloquinazoline scaffolds synthesized by a simple and highly selective Ugi/cyclization sequence. *Org Biomol Chem* 15:7549–7557. <https://doi.org/10.1039/c7ob01807j>
192. Pertejo P, Carreira-Barral I, Pena-Calleja P, Quesada R, Garcia-Valverde M (2020) Post-Ugi transformations for the access to pyrrolobenzodiazepine scaffolds with different degrees of unsaturation. *J Org Chem* 85:2291–2302. <https://doi.org/10.1021/acs.joc.9b02995>
193. Sañudo M, García-Valverde M, Marcaccini S, Delgado JJ, Rojo J, Torroba T (2009) Synthesis of benzodiazepine  $\beta$ -turn mimetics by an Ugi 4CC/Staudinger/aza-Wittig sequence. Solving the conformational behavior of the Ugi 4CC adducts. *J Org Chem* 74:2189–2192. <https://doi.org/10.1021/jo8025862>
194. Lecinska P, Corres N, Moreno D, García-Valverde M, Marcaccini S, Torroba T (2010) Synthesis of pseudo-peptidic (*S*)-6-amino-5-oxo-1,4-diazepines and (*S*)-3-benzyl-2-oxo-1,4-benzodiazepines by an Ugi 4CC Staudinger/aza-Wittig sequence. *Tetrahedron* 66:6783–6788. <https://doi.org/10.1016/j.tet.2010.06.062>
195. Corres N, Delgado JJ, García-Valverde M, Marcaccini S, Rodríguez T, Rojo J, Torroba T (2008) Synthesis of pseudo-peptidic [(*5H*)-6-oxodibenzo[*b*, *f*][1,5]diazocine-5-yl]arylglycinamides by an Ugi 4CC/Staudinger/aza-Wittig sequence. *Tetrahedron* 64:2225–2232. <https://doi.org/10.1016/j.tet.2007.12.028>
196. Pertejo P, Corres N, Torroba T, Garcia-Valverde M (2015) Reversal of diastereoselectivity in the synthesis of peptidomimetic 3-carboxamide-1,4-benzodiazepin-5-ones. *Org Lett* 17:612–615. <https://doi.org/10.1021/ol503628r>
197. Yan Y-M, Gao Y, Ding M-W (2016) New efficient synthesis of multisubstituted benzimidazoles and quinoxalin-2(*1H*)-ones by a Ugi 4CC/aza-Wittig sequence starting from aromatic amine precursors. *Tetrahedron* 72:5548–5557. <https://doi.org/10.1016/j.tet.2016.07.048>
198. Welsch SJ, Umkehrer M, Kalinski C, Ross G, Burdack C, Kolb J, Wild M, Ehrlich A, Wessjohann LA (2015) Synthesis of substituted imidazolines by an Ugi/Staudinger/aza-Wittig sequence. *Tetrahedron Lett* 56:1025–1029. <https://doi.org/10.1016/j.tetlet.2015.01.043>
199. Zhong Y, Wang L, Ding M-W (2011) New efficient synthesis of 2,3,4-trisubstituted 3,4-dihydroquinazolines by a Ugi 4CC/Staudinger/aza-Wittig sequence. *Tetrahedron* 67:3714–3723. <https://doi.org/10.1016/j.tet.2011.03.056>
200. Zhong Y, Zhang H, Ding M-W (2015) Regioselective synthesis of 2-acylquinazolines and 3*H*-1,4-benzodiazepin-3-ones by a Ugi 4CC/Staudinger/aza-Wittig sequence. *J Heterocycl Chem* 52:330–335. <https://doi.org/10.1002/jhet.1953>
201. He P, Nie Y-B, Wu J, Ding M-W (2011) Unexpected synthesis of indolo[1,2-*c*]quinazolines by a sequential Ugi 4CC–Staudinger–aza-Wittig–nucleophilic addition reaction. *Org Biomol Chem* 9:1429–1436. <https://doi.org/10.1039/C0OB00855A>
202. Ding M-W, Xie H, Liu J-C (2016) Facile synthesis of 3-arylidene-3*H*-1,4-benzodiazepines by a sequential Ugi/Staudinger/aza-Wittig reaction. *Synthesis* 48:4541–4547. <https://doi.org/10.1055/s-0036-1588308>
203. Ding M-W, Wang L, Ren Z-L, Chen M (2014) One-pot synthesis of 2,4,5-trisubstituted oxazoles via a tandem Passerini three-component coupling/Staudinger/aza-Wittig/isomerization reaction. *Synlett* 25:721–723. <https://doi.org/10.1055/s-0033-1340596>
204. Ding M-W, Ren Z-L, Liu J-C (2016) A facile synthesis of 4-tetra-*z*-substituted 4*H*-3,1-benzoxazines through sequential Passerini-azide/acylation/catalytic aza-Wittig reaction. *Synthesis* 49:745–754. <https://doi.org/10.1055/s-0036-1588333>
205. Akbarzadeh R, Amanpour T, Bazgir A (2014) Synthesis of 3-oxo-1,4-diazepine-5-carboxamides and 6-(4-oxo-chromen-3-yl)-pyrazinones via sequential Ugi 4CC/Staudinger/intramolecular nucleophilic cyclization and Ugi 4CC/Staudinger/aza-Wittig reactions. *Tetrahedron* 70:8142–8147. <https://doi.org/10.1016/j.tet.2014.07.102>
206. Marcaccini S, Pepino R, Polo C, Pozo C (2001) Studies on isocyanides and related compounds; a facile synthesis of 4-phenyl-1-(*2H*)phthalazinone-2-alkanoic acid amides. *Synthesis* 2001:85–88. <https://doi.org/10.3184/030823401103168695>
207. Sañudo M, Marcaccini S, Basurto S, Torroba T (2006) Synthesis of 3-hydroxy-6-oxo[1,2,4]triazin-1-yl alaninamides, a new class of cyclic dipeptidyl ureas. *J Org Chem* 71:4578–4584. <https://doi.org/10.1021/jo060434y>
208. Bossio R, Marcaccini S, Pepino R, Torroba T (1993) Studies on isocyanides and related compounds - a novel synthetic route to furan derivatives. *Synthesis* 1993:783–785. <https://doi.org/10.1055/s-1993-25940>
209. Bossio R, Marcaccini S, Pepino R (1994) Studies on isocyanides and related compounds - synthesis of a novel class of furan derivatives. *Liebigs Ann Chem* 1994:527–528. <https://doi.org/10.1002/jlac.15619940515>
210. Bossio R, Marcaccini S, Pepino R, Torroba T (1999) Studies on isocyanides and related compounds. A facile synthesis of 1-substituted 3-cyano-2-methoxy-3-phenylpyrroles. *Heterocycles* 50:463–467. [https://doi.org/10.3987/COM-98-S\(H\)16](https://doi.org/10.3987/COM-98-S(H)16)
211. Marcaccini S, Neo AG, Marcos CF (2009) Sequential five-component synthesis of spiropyrrolidinochromanones. *J Org Chem* 74:6888–6890. <https://doi.org/10.1021/jo900992w>
212. Marcaccini S, Pepino R, Paoli P, Rossi P, Torroba T (2001) Studies on isocyanides and related compounds. *N, N'*-disubstituted 2-cyanoacetoxy-2-cyanomethylmalondiamides from an unusual reaction between isocyanides and cyanoacetic acid. *J Chem Res Synop* 2001:465–467. <https://doi.org/10.3184/030823401103168695>
213. Bossio R, Marcaccini S, Pepino R (1991) Studies on isocyanides and related compounds - synthesis of oxazole derivatives via the

- Passerini reaction. *Liebigs Ann Chem* 1991:1107–1108. <https://doi.org/10.1002/jlac.1991199101193>
214. Bossio R, Marcaccini S, Pepino R, Polo C, Torroba T (1992) Synthesis of *N*-substituted 2-alkyl-4-aryloxazole-5-carboxamides. *Org Prep Proced Int* 24:188–191. <https://doi.org/10.1080/00304949209355696>
215. Bossio R, Marcaccini S, Pepino R (1991) Studies on isocyanides and related compounds - synthesis of *N*-substituted 4-aryl-2-arylthiomethyloxazole-5-carboxamides. *J Chem Res Synop* 1991:320–321
216. Faggi C, Garcia-Valverde M, Marcaccini S, Pepino R, Pozo MC (2003) Studies on isocyanides and related compounds: A facile synthesis of 1,6-dihydro-6-oxopyrazine-2-carboxylic acid derivatives via Ugi four-component condensation. *Synthesis* 2003:1553–1558. <https://doi.org/10.1055/s-2003-40525>
217. Neo AG, Marcos CF, Marcaccini S, Pepino R (2005) Studies on isocyanides. A facile synthesis of 4,5-dihydro-1,4-benzothiazepin-3(2*H*)-ones via post-condensation modifications of the Ugi reaction. *Tetrahedron Lett* 46:7977–7979. <https://doi.org/10.1016/j.tetlet.2005.09.071>
218. Neo AG, Delgado J, Polo C, Marcaccini S, Marcos CF (2005) A new synthesis of  $\beta$ -keto amides by reduction of Passerini adducts. *Tetrahedron Lett* 46:23–26. <https://doi.org/10.1016/j.tetlet.2004.11.041>
219. Carrillo RM, Neo AG, Lopez-Garcia L, Marcaccini S, Marcos CF (2006) Zinc catalysed ester solvolysis. Application to the synthesis of tartronic acid derivatives. *Green Chem* 8:787–789. <https://doi.org/10.1039/b607358a>
220. Neo AG, Carrillo RM, Delgado J, Marcaccini S, Marcos CF (2011) A multicomponent approach to the synthesis of 1,3-dicarbonylic compounds. *Mol Divers* 15:529–539. <https://doi.org/10.1007/s11030-010-9277-y>
221. Kobayashi K, Shirai Y, Fukamachi S, Konishi H (2010) Synthesis of 2,3-bis(arylamino)benzofurans and 2,3-bis(arylimino)-2,3-dihydrobenzofurans by a Lewis acid catalyzed reaction of 2-aryliminophenols with aryl isocyanides. *Synthesis* 2010:666–670. <https://doi.org/10.1055/s-0029-1218609>
222. Ramazani A, Mahyari AT, Rouhani M, Rezaei A (2009) A novel three-component reaction of a secondary amine and a 2-hydroxybenzaldehyde derivative with an isocyanide in the presence of silica gel: An efficient one-pot synthesis of benzo[*b*]furan derivatives. *Tetrahedron Lett* 50:5625–5627. <https://doi.org/10.1016/j.tetlet.2009.07.115>
223. Mitra S, Hota S, Chattopadhyay P (2010) Lewis acid catalyzed one-pot selective synthesis of aminobenzofurans and *N*-alkyl-2-aryl-2-(arylimino)acetamides: Product dependence on the nature of the aniline. *Synthesis* 2010:3899–3905. <https://doi.org/10.1055/s-0030-1258239>
224. Adib M, Mahdavi M, Bagherzadeh S, Zhu L-G, Rahimi-Nasrabadi M (2010) Reaction between anthranilic acids, salicylaldehydes and isocyanides in water: An efficient synthesis of 2-[[2-(alkylimino)-1-benzofuran-3-ylidene]amino]benzoic acids. *Tetrahedron Lett* 51:27–29. <https://doi.org/10.1016/j.tetlet.2009.05.017>
225. García-González MC, González-Zamora E, Santillan R, Domínguez O, Méndez-Stivalet JM, Farfán N (2009) Three-component synthesis of 2-imino-1,4-benzoxazines. *Tetrahedron* 65:5337–5342. <https://doi.org/10.1016/j.tet.2009.04.080>
226. Okitsu T, Nagase K, Nishio N, Wada A (2012) Tandem migration-carboalkoxylation of *o*-isocyanophenyl acetals leading to benzoxazoles. *Org Lett* 14:708–711. <https://doi.org/10.1021/ol203175a>
227. Soeta T, Shitaya S, Okuno T, Fujinami S, Ukaji Y (2016) Efficient synthesis of benzothiophenes by [4+1] cycloaddition of 2-mercaptobenzaldehyde derivatives with isocyanides. *Tetrahedron* 72:7901–7905. <https://doi.org/10.1016/j.tet.2016.09.054>
228. Shirri M, Faghihi Z, Oskouei HA, Heravi MM, Fazelzadeh S, Notash B (2016) The synthesis of iminothiophenone-fused quinolines and evaluation of their serendipitous reactions. *RSC Adv* 6:92235–92240. <https://doi.org/10.1039/c6ra11469e>
229. Bossio R, Marcaccini S, Pepino R, Torroba T (1996) Studies on isocyanides and related compounds: synthesis of benzo[*c*]thiophenes by way of acid-induced three-component reactions. *J Chem Soc Perkin Trans 1*(1996):229–230. <https://doi.org/10.1039/P19960000229>
230. El Kaïm L, Grimaud L (2010) Ugi-Smiles couplings: New entries to *N*-aryl carboxamide derivatives. *Mol Divers* 14:855–867. <https://doi.org/10.1007/s11030-009-9175-3>
231. El Kaim L, Grimaud L, Oble J (2005) Phenol Ugi-smiles systems: Strategies for the multicomponent *N*-arylation of primary amines with isocyanides, aldehydes, and phenols. *Angew Chem Int Ed* 44:7961–7964. <https://doi.org/10.1002/anie.200502636>
232. Castellano TG, Neo AG, Marcaccini S, Marcos CF (2012) Enols as feasible acid components in the Ugi condensation. *Org Lett* 14:6218–6221. <https://doi.org/10.1021/ol302976g>
233. Neo AG, Castellano TG, Marcos CF (2015) Enol-Ugi reaction of hydroxycoumarins: Straightforward synthesis of amino acid derived coumarin enamines. *Synthesis* 47:2431–2438. <https://doi.org/10.1055/s-0034-1380436>
234. Neo AG, Castellano TG, Marcos CF (2017) An easy synthesis of diversely functionalized 2*H*-chromenes and amido amines by an enol-Ugi reaction. *ARKIVOC* 2017:21–31. <https://doi.org/10.3998/ark.5550190.p009.775>
235. Aknin K, Gauriot M, Totobenazara J, Deguine N, Deprez-Poulain R, Deprez B, Charton J (2012) Squaric acid is a suitable building-block in 4*C*-Ugi reaction: Access to original bivalent compounds. *Tetrahedron Lett* 53:458–461. <https://doi.org/10.1016/j.tetlet.2011.11.077>
236. Mehrabi H, Saleki T (2015) The double four-component Ugi reaction between squaramic or squaramide acids with alkyl isocyanides, amines and aromatic aldehydes. *J Chem Res* 39:195–198. <https://doi.org/10.3184/174751915x14266899341593>
237. Massoudi A, Amini I, Ramazani A, Nasrabadi FZ, Ahmadi Y (2012) Four-component synthesis of 2-(*N*, *N*-dialkylamino)-2,4,6-cycloheptatrien-1-one derivatives from tropolone, an isocyanide, a primary amine and an aldehyde via Ugi-Smiles coupling reaction. *Bull Korean Chem Soc* 33:39–42. <https://doi.org/10.5012/bkcs.2012.33.1.39>
238. Rabêlo WF, Echemendía R (2019) Synthesis of novel 1,4 naphthoquinone-based molecules by an Ugi-type four-component reaction. *Synth Commun* 49:515–521. <https://doi.org/10.1080/00397911.2018.1551548>
239. Ramezanzpour S, Bigdeli Z, Alavijeh N, Rominger F (2017) Application of saccharin as an acidic partner in the Ugi reaction for the one-pot synthesis of 3-iminosaccharins. *Synlett* 28:1214–1218. <https://doi.org/10.1055/s-0036-1558962>
240. Neo AG, Marcos CF (2018) Selective synthesis of 3-substituted pyrrolidinones by enol-Passerini and anomalous enol-Passerini condensations. *Org Lett* 20:3875–3878. <https://doi.org/10.1021/acs.orglett.8b01462>
241. Bornadiego A, Neo AG, Marcos CF (2021) Synthesis of chromeno[3,4-*b*]piperazines by an enol-Ugi/reduction/cyclization sequence. *Molecules* 26:1287–1304. <https://doi.org/10.3390/molecules26051287>
242. Echemendía R, Da Silva GP, Kawamura MY, De La Torre AF, Corrêa AG, Ferreira MAB, Rivera DG, Paixão MW (2019) A stereoselective sequential organocascade and multicomponent approach for the preparation of tetrahydropyridines and chimeric derivatives. *Chem Commun* 55:286–289. <https://doi.org/10.1039/c8cc06871b>

243. Echemendia R, de La Torre AF, Monteiro JL, Pila M, Correa AG, Westermann B, Rivera DG, Paixao MW (2015) Highly stereoselective synthesis of natural-product-like hybrids by an organocatalytic/multicomponent reaction sequence. *Angew Chem Int Ed* 54:7621–7625. <https://doi.org/10.1002/anie.201412074>
244. van Leusen AM, Siderius H, Hoogenboom BE, van Leusen D (1972) A new and simple synthesis of the pyrrole ring system from Michael acceptors and tosylmethylisocyanides. *Tetrahedron Lett* 13:5337–5340. [https://doi.org/10.1016/S0040-4039\(01\)85244-8](https://doi.org/10.1016/S0040-4039(01)85244-8)
245. Barton DHR, Kervagoret J, Zard SZ (1990) A useful synthesis of pyrroles from nitroolefins. *Tetrahedron* 46:7587–7598. [https://doi.org/10.1016/S0040-4020\(01\)89069-4](https://doi.org/10.1016/S0040-4020(01)89069-4)
246. Ono N, Muratani E, Ogawa T (1991) Preparation of 2-formyl-4-nitropyrroles. *J Heterocycl Chem* 28:2053–2055. <https://doi.org/10.1002/jhet.5570280846>
247. Delvalle JL, Polo C, Torroba T, Marcaccini S (1995) A short synthesis of *C*-glycosyl pyrazoles and pyrroles. *J Heterocycl Chem* 32:899–901. <https://doi.org/10.1002/jhet.5570320337>
248. ten Have R, Leusink FR, van Leusen AM (1996) An efficient synthesis of substituted 3(4)-nitropyrroles from nitroalkenes and tosylmethyl isocyanides. *Synthesis* 1996:871–876. <https://doi.org/10.1055/s-1996-4306>
249. Krishna PR, Reddy VVR, Sharma GVM (2003) A versatile protocol for the synthesis of oxazole and 3-nitro/3-carbomethoxy pyrrole *C*-nucleosides using TOSMIC. *Synlett* 2003:1619–1622. <https://doi.org/10.1055/s-2003-41007>
250. Krishna PR, Reddy VVR, Srinivas R (2007) A new synthetic route to oxazole and pyrrole 2-deoxy-*C*-ribosides. *Tetrahedron* 63:9871–9880. <https://doi.org/10.1016/j.tet.2007.06.110>
251. Bossio R, Marcaccini S, Pepino R (1986) A novel class of nitrile ylide. *Tetrahedron Lett* 27:4643–4646. [https://doi.org/10.1016/S0040-4039\(00\)85027-3](https://doi.org/10.1016/S0040-4039(00)85027-3)
252. Bossio R, Marcaccini S, Paoli P, Pepino R, Polo C (1990) Studies on isonitriles and related compounds - synthesis of 1*H*-pyrrole and 1*H*-imidazole derivatives via 1,3-dipolar cycloaddition. *Heterocycles* 31:1855–1860. <https://doi.org/10.3987/COM-90-5529>
253. Ito Y, Kato H, Saegusa T (1982) A new approach for stereoselective synthesis of  $\gamma$ -butyrolactones. *J Org Chem* 47:741–743. <https://doi.org/10.1021/jo00343a030>
254. Chatani N, Oshita M, Tobisu M, Ishii Y, Murai S (2003) A GaCl<sub>3</sub>-catalyzed [4+1] cycloaddition of  $\alpha$ ,  $\beta$ -unsaturated carbonyl compounds and isocyanides leading to unsaturated  $\gamma$ -lactone derivatives. *J Am Chem Soc* 125:7812–7813. <https://doi.org/10.1021/ja035014u>
255. Yavari I, Shaabani A, Maghsoodlou MT (1997) On the reaction between alkyl isocyanides and 3-benzylidene-2,4-pentanedione. A convenient synthetic route to densely functionalized furans. *Monatsh Chem* 128:697–700
256. Quai M, Frattini S, Vendrame U, Mondoni M, Dossena S, Cereda E (2004) 5-hydroxy-2*H*-pyrrol-2-ones and not 2-aminofurans are the cycloaddition products between alkyl isocyanides and benzylidene-1,3-diketones. *Tetrahedron Lett* 45:1413–1416
257. Neo AG, Bornadiego A, Diaz J, Marcaccini S, Marcos CF (2013) Elusive 2-aminofuran Diels-Alder substrates for a straightforward synthesis of polysubstituted anilines. *Org Biomol Chem* 11:6546–6555. <https://doi.org/10.1039/c3ob41411f>
258. Bornadiego A, Díaz J, Marcos CF (2014) Synthesis of 4-aminoxanthenes by an uncatalyzed, multicomponent reaction. *Adv Synth Catal* 356:718–722. <https://doi.org/10.1002/adsc.201300750>
259. Bornadiego A, Díaz J, Marcos CF (2015) Regioselective tandem [4+1]-[4+2] synthesis of amino-substituted dihydroxanthenes and xanthenes. *J Org Chem* 80:6165–6172. <https://doi.org/10.1021/acs.joc.5b00658>
260. Bornadiego A, Díaz Álvarez J, Marcos FC (2019) Tandem synthesis of 4-aminoxanthenes is controlled by a water-assisted tautomerization: A general straightforward reaction. *Org Biomol Chem* 17:1410–1422. <https://doi.org/10.1039/C8OB02527D>
261. Teimouri MB, Mohammadnia S (2018) Reaction between alkyl isocyanides and 3-formylchromones in the presence of monocyclic unsaturated acyl compounds: competition between Friedel-Crafts acylation and Diels-Alder cycloaddition. *Tetrahedron* 74:1767–1775. <https://doi.org/10.1016/j.tet.2018.01.054>
262. Teimouri MB, Savadjani ZB, Shiri M, Bikas R, Naderi S (2022) Combining cycloaddition reactions for the one-pot synthesis of novel xanthoquinone dyes. *Dyes Pigm* 199:110106. <https://doi.org/10.1016/j.dyepig.2022.110106>
263. Bornadiego A, Díaz J, Marcos CF (2019) Direct multicomponent synthesis of benzocoumarins. *Peer J Organic Chem* 1:e1-14. <https://doi.org/10.7717/peerj-ochem.1>
264. Men Y, Hu Z, Dong J, Xu X, Tang B (2018) Formal [1+2+3] annulation: Domino access to carbazoles and indolocarbazole alkaloids. *Org Lett* 20:5348–5352. <https://doi.org/10.1021/acs.orglett.8b02266>
265. Dong J, Zhang D, Men Y, Zhang X, Hu Z, Xu X (2019) [1+2+3] annulation as a general access to indolo[3,2-*b*]carbazoles: Synthesis of malasseziazole *C*. *Org Lett* 21:166–169. <https://doi.org/10.1021/acs.orglett.8b03646>
266. Bornadiego A, Díaz J, Marcos CF (2019) Tandem synthesis of polycyclic isoindoles. *J Org Chem* 84:7426–7433. <https://doi.org/10.1021/acs.joc.9b00381>
267. Gelman M, Massarano T, Lavi R, Byk G (2018) A new multicomponent reaction MCR4 for the synthesis of analogs of staurosporine. *Curr Org Chem* 22:505–517. <https://doi.org/10.2174/1385272821666170817110101>
268. Hu Z, Men Y, Xu Z, Wu T, Xu X, Tang B (2020) A catalyst-free aqueous mediated multicomponent reaction of isocyanide: Expedient synthesis of polyfunctionalized cyclo[*b*]fused mono-, di- and tricarbazoles. *Organic Chem Front* 7:3720–3726. <https://doi.org/10.1039/d0qo01095b>
269. Hu Z, Dong J, Li Z, Yuan B, Wei R, Xu X (2018) Metal-free triple annulation of ene-yne-ketones with isocyanides: Domino access to furan-fused heterocycles via furoketenimine. *Org Lett* 20:6750–6754. <https://doi.org/10.1021/acs.orglett.8b02870>
270. Díaz-Álvarez J, Neo AG, Carrillo RM, Garrido L, Delgado J, Marcos CF (2008) Synthesis of furochromones by [4+1] cycloaddition of 3-formylchromone with isocyanides 1st Portuguese Young Chemist Meeting (1stPYChem). Instituto Superior Técnico de Lisboa, Lisbon (Portugal), p 79
271. Díaz Álvarez J, Carrillo RM, Garrido L, Neo AG, Delgado J, Jover A, Marcaccini S, Marcos CF (2009) Synthesis of fluorescent chemosensors based in chromone core 16th European Symposium on Organic Chemistry (ESOC 2009). Institute of Organic Chemistry and Biochemistry Academy of Sciences of the Czech Republic, Prague (Czech Republic ), p 170
272. Neo AG, Garrido L, Diaz J, Marcaccini S, Marcos CF (2012) Furo [3,4-*b*] chromones, and not pyrano [3,4-*b*] chromones, are obtained by the reaction of 3-formylchromones with isocyanides. *Synlett* 2012:2227–2230. <https://doi.org/10.1055/s-0032-1317032>
273. Panja SK, Maiti S, Banerjee S (1914) Bandyopadhyay C (2010) One-pot synthesis of pyrano[3,4-*b*]chromones from chromone-3-carbaldehyde. *Synlett* 2010:1909. <https://doi.org/10.1055/s-0030-1258496>
274. Panja SK, Maiti S, Drew MGB, Bandyopadhyay C (2011) One-pot three component reaction involving cyclohexyl isocyanide for

- the synthesis of furo[3,4-*b*]chromenes. *J Chem Res* 35:225–228. <https://doi.org/10.3184/174751911X13015945166625>
275. Teimouri MB (2011) Serendipitous stereoselective synthesis of brand-new fluorescent dyes: (1 *Z*)-3-(alkylimino)-1-[(chromone-3-yl)methylene]-1,3-dihydro-9*h*-furo[3,4-*b*]chromen-9-one-type fluorophores with blue fluorescence emission properties. *Tetrahedron* 67:1837–1843. <https://doi.org/10.1016/j.tet.2011.01.033>
276. Terzidis MA, Tsiaras VG, Drosos NM, Kasapidou PM, Stephanidou-Stephanatou J, Tsoleridis CA, Buth G, Kostakis GE (2014) Chromeno[2,3-*c*]pyrroles by one-pot multicomponent domino addition–amination reaction. *Tetrahedron Lett* 55:5601–5604. <https://doi.org/10.1016/j.tetlet.2014.08.048>
277. Teimouri MB, Zolfaghari F, Naderi S (2017) Furochromone-isatin conjugates via an uncatalyzed diastereoselective [4+1] cycloaddition/tautomerization/Friedel-Crafts hydroxyalkylation domino reaction. *Tetrahedron* 73:262–271. <https://doi.org/10.1016/j.tet.2016.12.010>
278. Teimouri MB, Mokhtare Z, Khavasi HR (2021) Uncatalyzed diastereoselective synthesis of alkyliminofurochromone-derived benzylmalononitriles via a three-component cascade reaction: Competition between Diels-Alder cycloaddition and Michael addition. *Org Biomol Chem* 19:2517–2525. <https://doi.org/10.1039/d0ob02540b>
279. Teimouri MB, Batebi E, Mohammadnia S, Khavasi HR (2021) Water-controlled selectivity switch in a multicomponent reaction: One-pot stereoselective synthesis of (acyloxymethylidene) chromonyl-furochromones and amido-(acyloxymethylidene) chromones. *Tetrahedron* 96:132374. <https://doi.org/10.1016/j.tet.2021.132374>
280. Neo AG, Diaz J, Marcaccini S, Marcos CF (2012) Conjugate addition of isocyanides to chromone 3-carboxylic acid: an efficient one-pot synthesis of chroman-4-one 2-carboxamides. *Org Biomol Chem* 10:3406–3416. <https://doi.org/10.1039/c2ob07011a>
281. Shaabani A, Soleimani E, Rezayan AH, Sarvary A, Khavasi HR (2008) Novel isocyanide-based four-component reaction: a facile synthesis of fully substituted 3,4-dihydrocoumarin derivatives. *Org Lett* 10:2581–2584. <https://doi.org/10.1021/o1800856e>

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.